



PSGL-1 Rabbit pAb

db8742 Package: 20μL 50μL 100μL

Product Name: PSGL-1 Rabbit pAb

Cat.No.: db8742

 $\textbf{Synonyms:} \ \mathsf{CLA}; \ \mathsf{CD162}; \ \mathsf{PSGL1}; \ \mathsf{PSGL-1}$

Application: WB, IHC, ICC/IF, FC, IP

Reactivity : Human

Host species : Rabbit

Background This gene encodes a glycoprotein that functions as a high affinity counter-receptor for the cell

adhesion molecules P-, E- and L- selectin expressed on myeloid cells and stimulated T

lymphocytes. As such, this protein plays a critical role in leukocyte trafficking during inflammation by

tethering of leukocytes to activated platelets or endothelia expressing selectins. This protein

requires two post-translational modifications, tyrosine sulfation and the addition of the sialyl Lewis \boldsymbol{x}

tetrasaccharide (sLex) to its O-linked glycans, for its high-affinity binding activity. Aberrant

expression of this gene and polymorphisms in this gene are associated with defects in the innate and adaptive immune response. Alternate splicing results in multiple transcript variants. [provided

by RefSeq, Apr 2011]

Immunogen Recombinant protein of human PSGL-1

Gene ID 6404

Swiss Prot Q14242

Synonyms CLA; CD162; PSGL1; PSGL-1

Reactivity Human

Application WB, IHC, ICC/IF, FC, IP

Recommended dilution WB: 1:1000

IHC: 1:200 ICC/IF: 1:20 FC: 1:100

IP: 1:20

Calculated MW 43 kDa

Observed MW 250 kDa

Host species Rabbit



For Research Use Only **Product Datasheet**

Clonality Polyclonal

Isotype IgG

Purity Affinity Purification

Conjugation Un-conjugated

Storage Stability Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium

azide and 0.05% BSA. Stable for 12 months from date of receipt.