

## Phospho-eIF4EBP1/eIF4EBP2/eIF4EBP3 (Thr45) Rabbit pAb

db1574

Package : 20μL 50μL 100μL

**Product Name** : Phospho-eIF4EBP1/eIF4EBP2/eIF4EBP3 (Thr45) Rabbit pAb**Cat.No.:** db1574**Synonyms** : BP-1; 4EBP1; 4E-BP1; PHAS-I**Application** : WB, IHC, ICC/IF, IP**Reactivity** : Human, Mouse**Host species** : Rabbit**Background**

This gene encodes one member of a family of translation repressor proteins. The protein directly interacts with eukaryotic translation initiation factor 4E (eIF4E), which is a limiting component of the multisubunit complex that recruits 40S ribosomal subunits to the 5' end of mRNAs. Interaction of this protein with eIF4E inhibits complex assembly and represses translation. This protein is phosphorylated in response to various signals including UV irradiation and insulin signaling, resulting in its dissociation from eIF4E and activation of mRNA translation. [provided by RefSeq, Jul 2008]

**Immunogen**

A synthetic phosphopeptide corresponding to residues surrounding Thr45 of human eIF4EBP1

**Gene ID**

1978

**Swiss Prot**

Q13541

**Synonyms**

BP-1; 4EBP1; 4E-BP1; PHAS-I

**Reactivity**

Human, Mouse

**Application**

WB, IHC, ICC/IF, IP

**Recommended dilution**

WB: 1:2000  
IHC: 1:50  
ICC/IF: 1:50  
IP: 1:20

**Calculated MW**

13 kDa

**Observed MW**

15-20 kDa

**Host species**

Rabbit

**Clonality**

Polyclonal

**Isotype**

IgG

<b>Purity</b>	Affinity Purification
<b>Conjugation</b>	Un-conjugated
<b>Storage Stability</b>	Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.