

## Phospho-PKA alpha/beta/gamma (catalytic subunit) (Thr197) Rabbit pAb

db3644

Package : 20μL 50μL 100μL

**Product Name** : Phospho-PKA alpha/beta/gamma (catalytic subunit) (Thr197) Rabbit pAb

**Cat.No.:** db3644

**Synonyms** : PKACA; PPNAD4

**Application** : WB, IHC

**Reactivity** : Human, Mouse, Rat

**Host species** : Rabbit

### Background

This gene encodes one of the catalytic subunits of protein kinase A, which exists as a tetrameric holoenzyme with two regulatory subunits and two catalytic subunits, in its inactive form. cAMP causes the dissociation of the inactive holoenzyme into a dimer of regulatory subunits bound to four cAMP and two free monomeric catalytic subunits. Four different regulatory subunits and three catalytic subunits have been identified in humans. cAMP-dependent phosphorylation of proteins by protein kinase A is important to many cellular processes, including differentiation, proliferation, and apoptosis. Constitutive activation of this gene caused either by somatic mutations, or genomic duplications of regions that include this gene, have been associated with hyperplasias and adenomas of the adrenal cortex and are linked to corticotropin-independent Cushing's syndrome. Alternative splicing results in multiple transcript variants encoding different isoforms. Tissue-specific isoforms that differ at the N-terminus have been described, and these isoforms may differ in the post-translational modifications that occur at the N-terminus of some isoforms. [provided by RefSeq, Jan 2015]

### Immunogen

A synthetic phosphopeptide corresponding to residues surrounding Tyr197 of human PKA alpha/beta/gamma (catalytic subunit)

### Gene ID

5566

### Swiss Prot

P17612

### Synonyms

PKACA; PPNAD4

### Reactivity

Human, Mouse, Rat

### Application

WB, IHC

### Recommended dilution

WB: 1:1000  
IHC: 1:20-1:50

### Calculated MW

41 kDa

### Observed MW

41 kDa

<b>Host species</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	IgG
<b>Purity</b>	Affinity Purification
<b>Conjugation</b>	Un-conjugated
<b>Storage Stability</b>	Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.