

## Phospho-p38 (Thr180) Rabbit pAb

db5672

Package : 20μL 50μL 100μL

**Product Name** : Phospho-p38 (Thr180) Rabbit pAb**Cat.No.:** db5672**Synonyms** : RK; p38; CSBP; EXIP; Mxi2; CSBP1; CSBP2; CSPB1; PRKM14; PRKM15; SAPK2A; p38ALPHA**Application** : WB, FC**Reactivity** : Human**Host species** : Rabbit**Background**

The protein encoded by this gene is a member of the MAP kinase family. MAP kinases act as an integration point for multiple biochemical signals, and are involved in a wide variety of cellular processes such as proliferation, differentiation, transcription regulation and development. This kinase is activated by various environmental stresses and proinflammatory cytokines. The activation requires its phosphorylation by MAP kinase kinases (MKKs), or its autophosphorylation triggered by the interaction of MAP3K7IP1/TAB1 protein with this kinase. The substrates of this kinase include transcription regulator ATF2, MEK2C, and MAX, cell cycle regulator CDC25B, and tumor suppressor p53, which suggest the roles of this kinase in stress related transcription and cell cycle regulation, as well as in genotoxic stress response. Four alternatively spliced transcript variants of this gene encoding distinct isoforms have been reported. [provided by RefSeq, Jul 2008]

**Immunogen**

A synthetic phosphopeptide corresponding to residues surrounding Thr180 of human p38

**Gene ID**

1432

**Swiss Prot**

Q16539

**Synonyms**

RK; p38; CSBP; EXIP; Mxi2; CSBP1; CSBP2; CSPB1; PRKM14; PRKM15; SAPK2A; p38ALPHA

**Reactivity**

Human

**Application**

WB, FC

**Recommended dilution**WB: 1:1000  
FC: 1:50**Calculated MW**

41 kDa

**Observed MW**

41 kDa

**Host species**

Rabbit

**Clonality**

Polyclonal

Isotype	IgG
Purity	Affinity Purification
Conjugation	Un-conjugated
Storage Stability	Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.