

For Research Use Only **Product Datasheet**

PCNA (2E1) Mouse mAb

db6199 Package : 50μL 100μL

Product Name: PCNA (2E1) Mouse mAb

Cat.No.: db6199

Synonyms: Cyclin; DNA polymerase delta auxiliary protein; HGCN8729; MGC8367; Mutagen-sensitive 209

protein;OTTHUMP00000030189;OTTHUMP00000030190;PCNA;Pcna/cyclin;PCNA HUMAN;PCNAR

;Polymerase delta accessory protein;Proliferating Cell Nuclear Antigen

Application: WB, IP, ICC/IF, IHC-Fr, IHC-P

Reactivity: Human, Mouse, Rat, Monkey, Hamster

Host species: Mouse

Background Auxiliary protein of DNA polymerase delta and is involved in the control of eukaryotic DNA

replication by increasing the polymerase's processibility during elongation of the leading

strand.Induces a robust stimulatory effect on the 3'-5' exonuclease and 3'-phosphodiesterase,but

not apurinic-apyrimidinic (AP) endonuclease, APEX2 activities. Has to be loaded onto DNA in order

to be able to stimulate APEX2.Plays a key role in DNA damage response (DDR) by being

conveniently positioned at the replication fork to coordinate DNA replication with DNA repair and

DNA damage tolerance pathways. Acts as a loading platform to recruit DDR proteins that allow

completion of DNA replication after DNA damage and promote postreplication

repair/Monoubiquitinated PCNA leads to recruitment of translesion (TLS) polymerases, while 'Lys-63'-linked polyubiquitination of PCNA is involved in error-free pathway and employs recombination

mechanisms to synthesize across the lesion.

Immunogen Purified recombinant human PCNA protein fragments expressed in E.coli

Gene ID 5111

Swiss Prot P12004

Synonyms Cyclin; DNA polymerase delta auxiliary protein; HGCN8729; MGC8367; Mutagen-sensitive 209

protein;OTTHUMP00000030189;OTTHUMP00000030190;PCNA;Pcna/cyclin;PCNA_HUMAN;PCI

delta accessory protein; Proliferating Cell Nuclear Antigen

Reactivity Human, Mouse, Rat, Monkey, Hamster

Application WB, IP, ICC/IF, IHC-Fr, IHC-P

Recommended dilution WB: 1:1000

IHC: 1:50-1:200 ICC/IF: 1:100-1:200

CC/IF. 1.100-1.20

IP: 1:20



For Research Use Only **Product Datasheet**

Calculated MW 29 kDa

Observed MW 36 kDa

Host species Mouse

Clonality Monoclonal

Clonality No. 2E1-G10-H10

Isotype IgG2b


Purity Affinity Purification

Conjugation Un-conjugated


Storage Stability Store at -20°C. Supplied in PBS, 50% Glycerol(pH 7.3), 0.02% sodium azide and 0.5% BSA.

Stable for 12 months from date of receipt.






Western blot analysis of PCNA in Hela, 3T3, COS7, C6 and CHO-K1 lysates using PCNA antibody.



Immunocytochemistry analysis of PCNA in HeLa cells using PCNA antibody.



Immunoprecipitation analysis of PCNA (2E1) in Hela lysates using PCNA antibody.



Immunohistochemistry analysis of paraffin-embedded Human colorectal carcinoma using PCNA antibody (2E1G10H10). High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.