



ZAP70 (4B2) Mouse mAb

db6211 Package : 50μL 100μL

Product Name: ZAP70 (4B2) Mouse mAb

Cat.No.: db6211

Synonyms: SRK; STD; TZK; STCD; ZAP-70

Application: WB, IP
Reactivity: Human
Host species: Mouse

Background

Tyrosine kinase that plays an essential role in regulation of the adaptive immune response.Regulates motility,adhesion and cytokine expression of mature T-cells,as well as thymocyte development. Contributes also to the development and activation of primary Blymphocytes. When antigen presenting cells (APC) activate T-cell receptor (TCR), a serie of phosphorylations lead to the recruitment of ZAP70 to the doubly phosphorylated TCR component CD247/CD3Z through ITAM motif at the plasma membrane. This recruitment serves to localization to the stimulated TCR and to relieve its autoinhibited conformation. Release of ZAP70 active conformation is further stabilized by phosphorylation mediated by LCK.Subsequently,ZAP70 phosphorylates at least 2 essential adapter proteins/ LAT and LCP2. In turn,a large number of signaling molecules are recruited and ultimately lead to lymphokine production, T-cell proliferation and differentiation. Furthermore, ZAP70 controls cytoskeleton modifications, adhesion and mobility of T-lymphocytes, thus ensuring correct delivery of effectors to the APC.ZAP70 is also required for TCR-CD247/CD3Z internalization and degradation through interaction with the E3 ubiquitin-protein ligase CBL and adapter proteins SLA and SLA2. Thus, ZAP70 regulates both T-cell activation switch on and switch off by modulating TCR expression at the T-cell surface. During thymocyte development, ZAP70 promotes survival and cell-cycle progression of developing thymocytes before positive selection (when cells are still CD4/CD8 double negative). Additionally, ZAP70-dependent signaling pathway may also contribute to primary B-cells formation and activation through B-cell receptor (BCR).

Immunogen Purified recombinant human ZAP-70 protein fragments expressed in E.coli

Gene ID 7535

Swiss Prot P43403

Synonyms SRK; STD; TZK; STCD; ZAP-70

Reactivity Human

Application WB. IP





Recommended dilution WB: 1:500-1:1000

IP: 1:20

Calculated MW 70 kDa

Observed MW 70 kDa

Host species Mouse

Clonality Monoclonal

Clonality No. 4B2-C1-H3


Isotype lgG2b

Purity Affinity Purification


Conjugation Un-conjugated

Storage Stability Store at -20°C. Supplied in PBS, 50% Glycerol(pH 7.3), 0.02% sodium azide and 0.5% BSA.

Stable for 12 months from date of receipt.



Western blot analysis of ZAP70 in CEM and Jurkat lysates, negative in the Ramos lysates using ZAP70 antibody.



Immunoprecipitation analysis of ZAP70 (4B2) in Jurkat lysates (ZAP70 positive expression cell line)and Ramos lysates(ZAP70 negative expression cell line) using ZAP70 antibody.