

**ERK1/2 (4A4) Mouse mAb**

db6122

Package : 50µL 100µL

**Product Name :** ERK1/2 (4A4) Mouse mAb**Cat.No.:** db6122**Synonyms :** ERK; p38; p40; p41; ERK2; ERT1; ERK-2; MAPK2; PRKM1; PRKM2; P42MAPK; p41mapk; p42-MAPK; ERK1; ERT2; ERK-1; PRKM3; P44ERK1; P44MAPK; HS44KDAP; HUMKER1A; p44-ERK1; p44-MAPK**Application :** WB**Reactivity :** Human, Mouse, Rat**Host species :** Mouse

<b>Background</b>	Serine/threonine kinase which acts as an essential component of the MAP kinase signal transduction pathway. MAPK1/ERK2 and MAPK3/ERK1 are the 2 MAPKs which play an important role in the MAPK/ERK cascade. They participate also in a signaling cascade initiated by activated KIT and KITLG/SCF. Depending on the cellular context, the MAPK/ERK cascade mediates diverse biological functions such as cell growth, adhesion, survival and differentiation through the regulation of transcription, translation, cytoskeletal rearrangements.
<b>Immunogen</b>	Human p44 MAPK (Erk1) synthetic peptide conjugated to KLH
<b>Gene ID</b>	5595, 5594
<b>Swiss Prot</b>	P27361, P28482
<b>Synonyms</b>	ERK; p38; p40; p41; ERK2; ERT1; ERK-2; MAPK2; PRKM1; PRKM2; P42MAPK; p41mapk; p42-MAPK; ERK1; ERT2; ERK-1; PRKM3; P44ERK1; P44MAPK; HS44KDAP; HUMKER1A; p44-ERK1; p44-MAPK
<b>Reactivity</b>	Human, Mouse, Rat
<b>Application</b>	WB
<b>Recommended dilution</b>	WB: 1:500-1:1000
<b>Calculated MW</b>	44,42 kDa
<b>Observed MW</b>	42,44 kDa
<b>Host species</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Clonality No.</b>	4A4-G4-F6
<b>Isotype</b>	IgG1

<b>Purity</b>	Affinity Purification
<b>Conjugation</b>	Un-conjugated
<b>Storage Stability</b>	Store at -20°C. Supplied in PBS, 50% Glycerol(pH 7.3), 0.02% sodium azide and 0.5% BSA . Stable for 12 months from date of receipt.

