

## LC3A/B (3E9) Mouse mAb

db6313

Package : 50µL 100µL

**Product Name** : LC3A/B (3E9) Mouse mAb**Cat.No.:** db6313**Synonyms** : LC3; LC3A; ATG8E; MAP1ALC3; MAP1BLC3; MAP1LC3A; LC3B; ATG8F; MAP1LC3B-a;  
MAP1A/1BLC3; MAP1LC3B**Application** : WB**Reactivity** : Human, Rat**Host species** : Mouse**Background**

Macroautophagy is the major inducible pathway for the general turnover of cytoplasmic constituents in eukaryotic cells, it is also responsible for the degradation of active cytoplasmic enzymes and organelles during nutrient starvation. Macroautophagy involves the formation of double-membrane bound autophagosomes which enclose the cytoplasmic constituent targeted for degradation in a membrane bound structure, which then fuse with the lysosome (or vacuole) releasing a single-membrane bound autophagic bodies which are then degraded within the lysosome (or vacuole). MAP1A and MAP1B are microtubule-associated proteins which mediate the physical interactions between microtubules and components of the cytoskeleton. These proteins are involved in formation of autophagosomal vacuoles (autophagosomes). MAP1A and MAP1B each consist of a heavy chain subunit and multiple light chain subunits. MAP1LC3a is one of the light chain subunits and can associate with either MAP1A or MAP1B. The precursor molecule is cleaved by APG4B/ATG4B to form the cytosolic form, LC3-I. This is activated by APG7L/ATG7, transferred to ATG3 and conjugated to phospholipid to form the membrane-bound form, LC3-II.

**Immunogen**

Synthetic peptide corresponding to human LC3B protein

**Gene ID**

84557, 81631

**Swiss Prot**

Q9H492, Q9GZQ8

**Synonyms**LC3; LC3A; ATG8E; MAP1ALC3; MAP1BLC3; MAP1LC3A; LC3B; ATG8F; MAP1LC3B-a;  
MAP1A/1BLC3; MAP1LC3B**Reactivity**

Human, Rat

**Application**

WB

**Recommended dilution**

WB: 1:500-1:1000


**Calculated MW**

14 kDa

**Observed MW**

14,16 kDa

Host species	Mouse
Clonality	Monoclonal
Clonality No.	3E9-E5-C9
Isotype	IgG2b
Purity	Affinity Purification
Conjugation	Un-conjugated
Storage Stability	Store at -20°C. Supplied in PBS, 50% Glycerol(pH 7.3), 0.02% sodium azide and 0.5% BSA . Stable for 12 months from date of receipt.



Western blot analysis of LC3A/B in rat Cerebral Cortex, rat Brain and Hela lysates using LC3A/B antibody.