

CD45 Mouse mAb

db6409

Package : 50μL 100μL

Product Name : CD45 Mouse mAb**Cat.No.:** db6409**Synonyms** : PTPRC; CD45; Receptor-type tyrosine-protein phosphatase C; Leukocyte common antigen; L-CA; T200; CD antigen CD45**Application** : WB, IHC-P**Reactivity** : Human, Mouse**Host species** : Mouse**Background**

The protein encoded by this gene is a member of the protein tyrosine phosphatase (PTP) family. PTPs are known to be signaling molecules that regulate a variety of cellular processes including cell growth, differentiation, mitosis, and oncogenic transformation. This PTP contains an extracellular domain, a single transmembrane segment and two tandem intracytoplasmic catalytic domains, and thus is classified as a receptor type PTP. This PTP has been shown to be an essential regulator of T- and B-cell antigen receptor signaling. It functions through either direct interaction with components of the antigen receptor complexes, or by activating various Src family kinases required for the antigen receptor signaling. This PTP also suppresses JAK kinases, and thus functions as a regulator of cytokine receptor signaling. Alternatively spliced transcripts variants of this gene, which encode distinct isoforms, have been reported.

Immunogen

Purified recombinant fragment of human PTPRC (AA/ 928-989) expressed in E. Coli

Gene ID

5788

Swiss Prot

P08575

Synonyms

PTPRC; CD45; Receptor-type tyrosine-protein phosphatase C; Leukocyte common antigen; L-CA; T200; CD antigen CD45

Reactivity

Human, Mouse

Application

WB, IHC-P

Recommended dilutionWB: 1:500-1:1000
IHC: 1:50-1:100**Calculated MW**

147 kDa

Observed MW

180-240 kDa


Host species

Mouse


Clonality

Monoclonal


Clonality No.	5D3A3
Isotype	IgG1
Purity	Affinity Purification
Conjugation	Un-conjugated
Storage Stability	Store at -20°C. Supplied in PBS, 50% Glycerol(pH 7.3), 0.02% sodium azide and 0.5% BSA . Stable for 12 months from date of receipt.



Immunohistochemistry analysis of paraffin-embedded breast cancer tissues using PTPRC antibody with DAB staining.High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.



Immunohistochemistry analysis of paraffin-embedded esophageal cancer tissues using PTPRC antibody with DAB staining.High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.



Western blot analysis of PTPRC antibody in Ramos, Raw264.7 lysates using PTPRC antibody.