
# NF-kB p65 (3D2) Mouse mAb

**db6456** **Package : 50μL 100μL**


**Product Name :** NF-kB p65 (3D2) Mouse mAb
**Cat.No.:** db6456
**Synonyms :** NFKB3; RELA; TF65; Transcription factor p65; p65; NFkB
**Application :** WB, IHC-Fr, IHC-P, ICC/IF, IP
**Reactivity :** Human, Mouse, Rat
**Host species :** Mouse

| | |
|---|---|
| **Background** | NF-kappa-B is a pleiotropic transcription factor present in almost all cell types and is the endpoint of a series of signal transduction events that are initiated by a vast array of stimuli related to many biological processes such as inflammation, immunity, differentiation, cell growth, tumorigenesis and apoptosis. NF-kappa-B is a homo- or heterodimeric complex formed by the Rel-like domain-containing proteins RELA/p65, RELB, NFKB1/p105, NFKB1/p50, REL and NFKB2/p52 and the heterodimeric p65-p50 complex appears to be most abundant one. The dimers bind at kappa-B sites in the DNA of their target genes and the individual dimers have distinct preferences for different kappa-B sites that they can bind with distinguishable affinity and specificity. Different dimer combinations act as transcriptional activators or repressors, respectively. NF-kappa-B is controlled by various mechanisms of post-translational modification and subcellular compartmentalization as well as by interactions with other cofactors or corepressors. NF-kappa-B complexes are held in the cytoplasm in an inactive state complexed with members of the NF-kappa-B inhibitor (I-kappa-B) family. In a conventional activation pathway, I-kappa-B is phosphorylated by I-kappa-B kinases (IKKs) in response to different activators, subsequently degraded thus liberating the active NF-kappa-B complex which translocates to the nucleus. NF-kappa-B heterodimeric p65-p50 and p65-c-Rel complexes are transcriptional activators. The NF-kappa-B p65-p65 complex appears to be involved in invasin-mediated activation of IL-8 expression. The inhibitory effect of I-kappa-B upon NF-kappa-B the cytoplasm is exerted primarily through the interaction with p65. p65 shows a weak DNA-binding site which could contribute directly to DNA binding in the NF-kappa-B complex. Associates with chromatin at the NF-kappa-B promoter region via association with DDX1. Essential for cytokine gene expression in T-cells (PubMed/15790681). |
| **Immunogen** | Recombinant Protein of Transcription factor p65 |
| **Gene ID** | 5970 |
| **Swiss Prot** | Q04206 |
| **Synonyms** | NFKB3; RELA; TF65; Transcription factor p65; p65; NFkB |


| | |
|---|---|
| **Reactivity** | Human, Mouse, Rat |
| **Application** | WB, IHC-Fr, IHC-P, ICC/IF, IP |
| **Recommended dilution** | WB: 1:500-1000<br>IP: 1:20<br>ICC/IF: 1:50-1:200<br>IHC: 1:50-100 |
| **Calculated MW** | 60 kDa |
| **Observed MW** | 65 kDa |
| **Host species** | Mouse |
| **Clonality** | Monoclonal |
| **Clonality No.** | 3D2-4E9-7A8 |
| **Isotype** | IgG1 |
| **Purity** | Affinity Purification |
| **Conjugation** | Un-conjugated |
| **Storage Stability** | Store at -20°C. Supplied in PBS, 50% Glycerol(pH 7.3), 0.02% sodium azide and 0.5% BSA .<br>Stable for 12 months from date of receipt. |





Western blot analysis of NF-KB p65 in T47D, 3T3, COS7, C6 and Hela lysates using NF-KB p65 antibody.




Immunofluorescence analysis of NF-KB p65 (3D2) in Hela using NF-KB p65 (3D2) antibody,and DAPI (blue).




Immunoprecipitation analysis of NF-KB p65 (3D2) in Hela lysates using NFκB p65 (3D2) antibody.



Immunohistochemistry analysis of paraffin-embedded rat Brain Tissue using NF-KB p65 antibody.High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.




Western blot analysis of NF-KB p65 (3D2) in COS7, C6, T47D lysates using NF-KB p65 (3D2) antibody.