

## Caspase-8 (4C2) Mouse mAb

db6486

Package : 50µL 100µL

**Product Name** : Caspase-8 (4C2) Mouse mAb**Cat.No.:** db6486**Synonyms** : CASP8; MCH5; Caspase-8; CASP-8; Apoptotic cysteine protease; Apoptotic protease Mch-5; CAP4; FADD-homologous ICE/ced-3-like protease; FADD-like ICE; FLICE; ICE-like apoptotic protease 5; MORT1-associated ced-3 homolog; MACH**Application** : WB, ICC/IF, IHC-P**Reactivity** : Human, Mouse, Rat**Host species** : Mouse**Background**

Most upstream protease of the activation cascade of caspases responsible for the TNFRSF6/FAS mediated and TNFRSF1A induced cell death. Binding to the adapter molecule FADD recruits it to either receptor. The resulting aggregate called death-inducing signaling complex (DISC) performs CASP8 proteolytic activation. The active dimeric enzyme is then liberated from the DISC and free to activate downstream apoptotic proteases. Proteolytic fragments of the N-terminal propeptide (termed CAP3, CAP5 and CAP6) are likely retained in the DISC. Cleaves and activates CASP3, CASP4, CASP6, CASP7, CASP9 and CASP10. May participate in the GZMB apoptotic pathways. Cleaves ADPRT. Hydrolyzes the small-molecule substrate, Ac-Asp-Glu-Val-Asp-[AMC. Likely target for the cowpox virus CRMA death inhibitory protein. Isoform 5, isoform 6, isoform 7 and isoform 8 lack the catalytic site and may interfere with the pro-apoptotic activity of the complex.

**Immunogen**

Recombinant Protein of Caspase-8

**Gene ID**

841

**Swiss Prot**

Q14790

**Synonyms**

CASP8; MCH5; Caspase-8; CASP-8; Apoptotic cysteine protease; Apoptotic protease Mch-5; CAP4; FADD-homologous ICE/ced-3-like protease; FADD-like ICE; FLICE; ICE-like apoptotic protease 5; MORT1-associated ced-3 homolog; MACH

**Reactivity**

Human, Mouse, Rat

**Application**

WB, ICC/IF, IHC-P

**Recommended dilution**

WB: 1:500-1:1000

IHC: 1:50-1:100

ICC/IF: 1:50-1:200

**Calculated MW**

55 kDa

**Observed MW**

43,57 kDa

Host species	Mouse
Clonality	Monoclonal
Clonality No.	4C2-3G4-5G7
Isotype	IgG1
Purity	Affinity Purification
Conjugation	Un-conjugated
Storage Stability	<p>Store at -20°C. Supplied in PBS, 50% Glycerol(pH 7.3), 0.02% sodium azide and 0.5% BSA .</p> <p>Stable for 12 months from date of receipt.</p> <ul style="list-style-type: none"><li>□ Western blot analysis of Caspase8 in K562, 3T3, MCF-7 and Jurkat lysates using Caspase8 antibody.</li><li>□ Immunohistochemistry analysis of paraffin-embedded mouse Spleen Tissue using Caspase8 antibody.High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.</li><li>□ Western blot analysis of Caspase 8 (2G12) in THP-1, rat Brain, Hela lysates using Caspase8 (4C2) antibody.</li><li>□ Western blot analysis of Caspase 8 (2G12) in Jurkat, MOLT4 lysates using Caspase 8 (2G12) antibody</li><li>□ Immunocytochemistry analysis of Caspase8 (4C2) (green) in Hela using Caspase8 (4C2) antibody,and DAPI(blue).</li></ul>