

HDAC1 (3F5) Mouse mAb

db6538

Package : 50µL 100µL

Product Name : HDAC1 (3F5) Mouse mAb**Cat.No.:** db6538**Synonyms** : HDAC1; RPD3L1; Histone deacetylase 1; HD1**Application** : IHC-P**Reactivity** : Human, Rat, Mouse**Host species** : Mouse**Background**

Responsible for the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4). Histone deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events. Histone deacetylases act via the formation of large multiprotein complexes. Deacetylates SP proteins, SP1 and SP3, and regulates their function. Component of the BRG1-RB1-HDAC1 complex, which negatively regulates the CREB-mediated transcription in resting neurons. Upon calcium stimulation, HDAC1 is released from the complex and CREBBP is recruited, which facilitates transcriptional activation. Deacetylates TSHZ3 and regulates its transcriptional repressor activity. Deacetylates 'Lys-310' in RELA and thereby inhibits the transcriptional activity of NF-kappa-B. Deacetylates NR1D2 and abrogates the effect of KAT5-mediated relieving of NR1D2 transcription repression activity. Component of a RCOR/GFI/KDM1A/HDAC complex that suppresses, via histone deacetylase (HDAC) recruitment, a number of genes implicated in multilineage blood cell development. Involved in CIART-mediated transcriptional repression of the circadian transcriptional activator/ CLOCK-ARNTL/BMAL1 heterodimer. Required for the transcriptional repression of circadian target genes, such as PER1, mediated by the large PER complex or CRY1 through histone deacetylation.

Immunogen

Synthetic peptide conjugated to KLH

Gene ID

3065

Swiss Prot

Q13547

Synonyms

HDAC1; RPD3L1; Histone deacetylase 1; HD1

Reactivity

Human, Rat, Mouse

Application

IHC-P


Recommended dilution

IHC: 1:50-1:100


Calculated MW

62 kDa

Host species	Mouse
Clonality	Monoclonal
Clonality No.	3F5-2E8-1F7
Isotype	IgG1
Purity	Affinity Purification
Conjugation	Un-conjugated
Storage Stability	Store at -20°C. Supplied in PBS, 50% Glycerol(pH 7.3), 0.02% sodium azide and 0.5% BSA . Stable for 12 months from date of receipt.



Immunohistochemistry analysis of paraffin-embedded Human Colon Carcinoma Tissue using HDAC1 (3F5) antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.



Immunohistochemical analysis of paraffin-embedded Human tonsils using HDAC1 (3F5) antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.