
Caspase-3 (1B10) Mouse mAb

db6540 **Package : 50μL 100μL**


Product Name : Caspase-3 (1B10) Mouse mAb
Cat.No.: db6540
Synonyms : Caspase-3; CASP-3; Apopain; Cysteine protease CPP32; CPP-32; Protein Yama; SREBP cleavage activity 1; SCA-1; Caspase-3 subunit p17; Caspase-3 subunit p12; CASP3; CPP32
Application : IHC-P
Reactivity : Human, Rat, Mouse
Host species : Mouse

Background	Involved in the activation cascade of caspases responsible for apoptosis execution. At the onset of apoptosis it proteolytically cleaves poly(ADP-ribose) polymerase (PARP) at a '216-Asp-|-Gly-217' bond. Cleaves and activates sterol regulatory element binding proteins (SREBPs) between the basic helix-loop-helix leucine zipper domain and the membrane attachment domain. Cleaves and activates caspase-6, -7 and -9. Involved in the cleavage of huntingtin. Triggers cell adhesion in sympathetic neurons through RET cleavage.
Immunogen	Synthetic peptide conjugated to KLH
Gene ID	836
Swiss Prot	P42574
Synonyms	Caspase-3; CASP-3; Apopain; Cysteine protease CPP32; CPP-32; Protein Yama; SREBP cleavage activity 1; SCA-1; Caspase-3 subunit p17; Caspase-3 subunit p12; CASP3; CPP32
Reactivity	Human, Rat, Mouse
Application	IHC-P
Recommended dilution	IHC: 1:50-1:100
Host species	Mouse
Clonality	Monoclonal
Clonality No.	1B10-10E2-6A5
Isotype	IgG1
Purity	Affinity Purification
Conjugation	Un-conjugated
Storage Stability	Store at -20°C. Supplied in PBS, 50% Glycerol(pH 7.3), 0.02% sodium azide and 0.5% BSA .


Stable for 12 months from date of receipt.



Immunohistochemistry analysis of paraffin-embedded Human Skin Tissue using Caspase 3 (1B10) antibody.High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.



Immunohistochemical analysis of paraffin-embedded Human tonsils using Caspase 3 (1B10) antibody.High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.



Immunohistochemistry analysis of paraffin-embedded mouse Liver Tissue using Caspase3 antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.