

## ERK1 (1D10) Mouse mAb

db6569

Package : 50µL 100µL

**Product Name** : ERK1 (1D10) Mouse mAb**Cat.No.:** db6569**Synonyms** : MAPK; Mitogen-activated protein Kinase; PRKM2;MK01**Application** : IHC-P**Reactivity** : Human, Rat, Mouse**Host species** : Mouse**Background**

Serine/threonine kinase which acts as an essential component of the MAP kinase signal transduction pathway. MAPK1/ERK2 and MAPK3/ERK1 are the 2 MAPKs which play an important role in the MAPK/ERK cascade. They participate also in a signaling cascade initiated by activated KIT and KITLG/SCF. Depending on the cellular context, the MAPK/ERK cascade mediates diverse biological functions such as cell growth, adhesion, survival and differentiation through the regulation of transcription, translation, cytoskeletal rearrangements. The MAPK/ERK cascade plays also a role in initiation and regulation of meiosis, mitosis, and postmitotic functions in differentiated cells by phosphorylating a number of transcription factors. About 160 substrates have already been discovered for ERKs. Many of these substrates are localized in the nucleus, and seem to participate in the regulation of transcription upon stimulation. However, other substrates are found in the cytosol as well as in other cellular organelles, and those are responsible for processes such as translation, mitosis and apoptosis. Moreover, the MAPK/ERK cascade is also involved in the regulation of the endosomal dynamics, including lysosome processing and endosome cycling through the perinuclear recycling compartment (PNRC); as well as in the fragmentation of the Golgi apparatus during mitosis. The substrates include transcription factors (such as ATF2, BCL6, ELK1, ERF, FOS, HSF4 or SPZ1), cytoskeletal elements (such as CANX, CTTN, GJA1, MAP2, MAPT, PXN, SORBS3 or STMN1), regulators of apoptosis (such as BAD, BTG2, CASP9, DAPK1, IER3, MCL1 or PPARG), regulators of translation (such as EIF4EBP1) and a variety of other signaling-related molecules (like ARHGEF2, FRS2 or GRB10). Protein kinases (such as RAF1, RPS6KA1/RSK1, RPS6KA3/RSK2, RPS6KA2/RSK3, RPS6KA6/RSK4, SYK, MKNK1/MNK1, MKNK2/MNK2, RPS6KA5/MSK1, RPS6KA4/MSK2, MAPKAPK3 or MAPKAPK5) and phosphatases (such as DUSP1, DUSP4, DUSP6 or DUSP16) are other substrates which enable the propagation the MAPK/ERK signal to additional cytosolic and nuclear targets, thereby extending the specificity of the cascade.

**Immunogen**

Purified recombinant protein expressed in E.coli


**Gene ID**

5595


**Swiss Prot**

P27361

<b>Synonyms</b>	MAPK; Mitogen-activated protein Kinase; PRKM2;MK01
<b>Reactivity</b>	Human, Rat, Mouse
<b>Application</b>	IHC-P
<b>Recommended dilution</b>	IHC: 1:50-1:100
<b>Calculated MW</b>	44 kDa
<b>Host species</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Clonality No.</b>	1D10-8A3-8C6
<b>Isotype</b>	IgG1
<b>Purity</b>	Affinity Purification
<b>Conjugation</b>	Un-conjugated
<b>Storage Stability</b>	Store at -20°C. Supplied in PBS, 50% Glycerol(pH 7.3), 0.02% sodium azide and 0.5% BSA . Stable for 12 months from date of receipt.



Immunohistochemistry analysis of paraffin-embedded Human Tonsil Tissue using ERK1 (1D10) antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.



Immunohistochemical analysis of paraffin-embedded Human tonsils using ERK1 (1D10) antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.