

## ERK1/2 (1B4) Mouse mAb

db6570

Package : 50µL 100µL

**Product Name** : ERK1/2 (1B4) Mouse mAb**Cat.No.:** db6570**Synonyms** : MAPK3; ERK1; ERT2; ERK-1; PRKM3; P44ERK1; P44MAPK; HS44KDAP; HUMKER1A; p44-ERK1; p44-MAPK; MAPK1; ERK; p38; p40; p41; ERK2; ERT1; ERK-2; MAPK2; PRKM1; PRKM2; P42MAPK; p41mapk; p42-MAPK**Application** : WB, IHC-P**Reactivity** : Human, Rat, Mouse**Host species** : Mouse**Background**

erine/threonine kinase which acts as an essential component of the MAP kinase signal transduction pathway. MAPK1/ERK2 and MAPK3/ERK1 are the 2 MAPKs which play an important role in the MAPK/ERK cascade. They participate also in a signaling cascade initiated by activated KIT and KITLG/SCF. Depending on the cellular context, the MAPK/ERK cascade mediates diverse biological functions such as cell growth, adhesion, survival and differentiation through the regulation of transcription, translation, cytoskeletal rearrangements.

**Immunogen**

Synthetic peptide conjugated to KLH

**Gene ID**

5594, 5595

**Swiss Prot**

P27361, P28482

**Synonyms**

MAPK3; ERK1; ERT2; ERK-1; PRKM3; P44ERK1; P44MAPK; HS44KDAP; HUMKER1A; p44-ERK1; p44-MAPK; MAPK1; ERK; p38; p40; p41; ERK2; ERT1; ERK-2; MAPK2; PRKM1; PRKM2; P42MAPK; p41mapk; p42-MAPK

**Reactivity**

Human, Rat, Mouse

**Application**

WB, IHC-P

**Recommended dilution**WB: 1:500-1000  
IHC: 1:50-100**Calculated MW**

44,42 kDa

**Observed MW**

44,42 kDa

**Host species**

Mouse


**Clonality**

Monoclonal


**Clonality No.**

1B4-6A3-10A5


Isotype	IgG1
Purity	Affinity Purification
Conjugation	Un-conjugated
Storage Stability	Store at -20°C. Supplied in PBS, 50% Glycerol(pH 7.3), 0.02% sodium azide and 0.5% BSA . Stable for 12 months from date of receipt.



Western blot analysis of ERK1/2 (1B4)in Hela lysates, mouse Brain lysates, rat Brain lysates using ERK1/2 antibody.



Immunohistochemistry analysis of paraffin-embedded Human Lung Carcinoma Tissue using ERK1/2 antibody.High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.



Immunohistochemistry analysis of paraffin-embedded Human Colon Carcinoma Tissue using ERK1/2 antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.