

ATG7 (1F8) Mouse mAb**db6594****Package : 50µL 100µL****Product Name :** ATG7 (1F8) Mouse mAb**Cat.No.:** db6594**Synonyms :** hAGP7; Ubiquitin-activating enzyme E1-like protein; APG7L**Application :** IHC-P**Reactivity :** Human, Rat, Mouse**Host species :** Mouse**Background**

E1-like activating enzyme involved in the 2 ubiquitin-like systems required for cytoplasm to vacuole transport (Cvt) and autophagy. Activates ATG12 for its conjugation with ATG5 as well as the ATG8 family proteins for their conjugation with phosphatidylethanolamine. Both systems are needed for the ATG8 association to Cvt vesicles and autophagosomes membranes. Required for autophagic death induced by caspase-8 inhibition. Required for mitophagy which contributes to regulate mitochondrial quantity and quality by eliminating the mitochondria to a basal level to fulfill cellular energy requirements and preventing excess ROS production. Modulates p53/TP53 activity to regulate cell cycle and survival during metabolic stress. Plays also a key role in the maintenance of axonal homeostasis, the prevention of axonal degeneration, the maintenance of hematopoietic stem cells, the formation of Paneth cell granules, as well as in adipose differentiation.

Immunogen

Purified recombinant protein expressed in E.coli

Gene ID

10533

Swiss Prot

O95352

Synonyms

hAGP7; Ubiquitin-activating enzyme E1-like protein; APG7L

Reactivity

Human, Rat, Mouse

Application

IHC-P

Recommended dilution

IHC: 1:50-1:100

Calculated MW

78 kDa

Host species

Mouse

Clonality

Monoclonal


Clonality No.

1F8-8H4-4G10

Isotype

IgG1

Purity	Affinity Purification
Conjugation	Un-conjugated
Storage Stability	Store at -20°C. Supplied in PBS, 50% Glycerol(pH 7.3), 0.02% sodium azide and 0.5% BSA . Stable for 12 months from date of receipt.



Immunohistochemistry analysis of paraffin-embedded Human Breast Carcinoma Tissue using ATG7 (1F8) antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.