

STAT3 (6H10) Mouse mAb

db6609

Package : 50µL 100µL

Product Name : STAT3 (6H10) Mouse mAb**Cat.No.:** db6609**Synonyms** : STAT3; APRF; Signal transducer and activator of transcription 3; Acute-phase response factor**Application** : WB, IHC-P**Reactivity** : Human, Rat, Mouse**Host species** : Mouse**Background**

Signal transducer and transcription activator that mediates cellular responses to interleukins, KITLG/SCF, LEP and other growth factors (PubMed/10688651, PubMed/12359225, PubMed/12873986, PubMed/15194700, PubMed/17344214, PubMed/18242580, PubMed/23084476). Once activated, recruits coactivators, such as NCOA1 or MED1, to the promoter region of the target gene (PubMed/17344214). May mediate cellular responses to activated FGFR1, FGFR2, FGFR3 and FGFR4 (PubMed/12873986). Binds to the interleukin-6 (IL-6)-responsive elements identified in the promoters of various acute-phase protein genes (PubMed/12359225). Activated by IL31 through IL31RA (PubMed/15194700). Acts as a regulator of inflammatory response by regulating differentiation of naive CD4+ T-cells into T-helper Th17 or regulatory T-cells (Treg)/ deacetylation and oxidation of lysine residues by LOXL3, leads to disrupt STAT3 dimerization and inhibit its transcription activity (PubMed/28065600). Involved in cell cycle regulation by inducing the expression of key genes for the progression from G1 to S phase, such as CCND1 (PubMed/17344214). Mediates the effects of LEP on melanocortin production, body energy homeostasis and lactation . May play an apoptotic role by transactivating BIRC5 expression under LEP activation (PubMed/18242580). Cytoplasmic STAT3 represses macroautophagy by inhibiting EIF2AK2/PKR activity (PubMed/23084476). Plays a crucial role in basal beta cell functions, such as regulation of insulin secretion .

Immunogen

Purified recombinant protein expressed in E.coli

Gene ID

6774

Swiss Prot

P40763

Synonyms

STAT3; APRF; Signal transducer and activator of transcription 3; Acute-phase response factor

Reactivity

Human, Rat, Mouse

Application


WB, IHC-P

Recommended dilution


WB: 1:500-1:1000

IHC: 1:50-1:100


Calculated MW	88 kDa
Observed MW	79,86 kDa
Host species	Mouse
Clonality	Monoclonal
Clonality No.	6H10-4G1-4C9
Isotype	IgG1
Purity	Affinity Purification
Conjugation	Un-conjugated
Storage Stability	Store at -20°C. Supplied in PBS, 50% Glycerol(pH 7.3), 0.02% sodium azide and 0.5% BSA . Stable for 12 months from date of receipt.



Western blot analysis of STAT3 (6H10)in HeLa, 3T3, PC-12 lysates using STAT3 antibody.



Immunohistochemistry analysis of paraffin-embedded Human Colon Carcinoma using STAT3 antibody.High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.



Western blot analysis of STAT3 (6H10) in Jurkat, Raw264.7, Ramos, A549 lysates using Stat3 (6H1) antibody.