

## Histone H3 Rabbit pAb (HRP)

db6641

Package : 100μL 1mL

**Product Name** : Histone H3 Rabbit pAb (HRP)**Cat.No.:** db6641**Synonyms** : H3/A; H3FA**Application** : WB**Reactivity** : Human, Mouse, Rat**Host species** : Rabbit**Background**

Modulation of chromatin structure plays an important role in the regulation of transcription in eukaryotes. The nucleosome, made up of DNA wound around eight core histone proteins (two each of H2A, H2B, H3, and H4), is the primary building block of chromatin. The amino-terminal tails of core histones undergo various post-translational modifications, including acetylation, phosphorylation, methylation, and ubiquitination. These modifications occur in response to various stimuli and have a direct effect on the accessibility of chromatin to transcription factors and, therefore, gene expression. In most species, histone H2B is primarily acetylated at Lys5, 12, 15, and 20. Histone H3 is primarily acetylated at Lys9, 14, 18, 23, 27, and 56. Acetylation of H3 at Lys9 appears to have a dominant role in histone deposition and chromatin assembly in some organisms. Phosphorylation at Ser10, Ser28, and Thr11 of histone H3 is tightly correlated with chromosome condensation during both mitosis and meiosis (8-10). Phosphorylation at Thr3 of histone H3 is highly conserved among many species and is catalyzed by the kinase haspin. Immunostaining with phospho-specific antibodies in mammalian cells reveals mitotic phosphorylation at Thr3 of H3 in prophase and its dephosphorylation during anaphase.

**Immunogen**

A synthetic peptide corresponding to the carboxy terminus of the human histone H3 protein

**Gene ID**

8350

**Swiss Prot**

P68431

**Synonyms**

H3/A; H3FA

**Reactivity**

Human, Mouse, Rat

**Application**

WB

**Recommended dilution**

WB: 1:5000-1:10000

**Calculated MW**

15 kDa


**Observed MW**

15 kDa

**Host species**

Rabbit

Clonality	Polyclonal
Isotype	IgG
Purity	Affinity Purification
Conjugation	HRP
Storage Stability	Store at -20°C. Supplied in PBS, 50% Glycerol(pH 7.3), 0.02% sodium azide and 0.5% BSA . Stable for 12 months from date of receipt.



Western blot analysis of Histone H3 in various cell lines lysates using Histone H3 antibody(HRP Conjugate).