

Recombinant

DGRmAb®

## STAT5 (DGR11379) Rabbit mAb

db11277

Package : 10µL 20µL 50µL 100µL

**Product Name :** STAT5 (DGR11379) Rabbit mAb**Cat.No.:** db11277**Synonyms :** STAT5**Application :** WB, IHC-P, ICC/IF, FC**Reactivity :** Human,Mouse,Rat**Host species :** Rabbit**Background**

The protein encoded by this gene is a member of the STAT family of transcription factors. In response to cytokines and growth factors, STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. This protein mediates the signal transduction triggered by various cell ligands, such as IL2, IL4, CSF1, and different growth hormones. It has been shown to be involved in diverse biological processes, such as TCR signaling, apoptosis, adult mammary gland development, and sexual dimorphism of liver gene expression. This gene was found to fuse to retinoic acid receptor-alpha (RARA) gene in a small subset of acute promyelocytic leukemias (APLL). The dysregulation of the signaling pathways mediated by this protein may be the cause of the APLL. [provided by RefSeq, Jul 2008]

**Immunogen**

Recombinant protein of human STAT5B

**Gene ID**

6777

**Swiss Prot**

P51692

**Synonyms**

STAT5

**Reactivity**

Human,Mouse,Rat

**Application**

WB, IHC-P, ICC/IF, FC

**Recommended dilution**

WB: 1:1000-1:5000

IHC-P: 1:100

ICC/IF: 1:200-1:500

FC: 1:50-1:100

**Calculated MW**

90 kDa


**Observed MW**

90 kDa

**Host species**

Rabbit

Clonality	Monoclonal
Clonality No.	DGR11379
Isotype	IgG
Purity	Affinity Purification
Conjugation	Un-conjugated
Storage Stability	Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.



Western blot detection of STAT5B in C6 cell lysates using STAT5B antibody(1:1000 diluted).