

Recombinant

DGRmAb®

Bim (DGR11947) Rabbit mAb

db13581

Package : 10µL 20µL 50µL 100µL

Product Name : Bim (DGR11947) Rabbit mAb**Cat.No.:** db13581**Synonyms** : BAM; BIM; BOD**Application** : WB, IHC-P, ICC/IF, FC, IP**Reactivity** : Human,Mouse,Rat**Host species** : Rabbit**Background**

The protein encoded by this gene belongs to the BCL-2 protein family. BCL-2 family members form hetero- or homodimers and act as anti- or pro-apoptotic regulators that are involved in a wide variety of cellular activities. The protein encoded by this gene contains a Bcl-2 homology domain 3 (BH3). It has been shown to interact with other members of the BCL-2 protein family and to act as an apoptotic activator. The expression of this gene can be induced by nerve growth factor (NGF), as well as by the forkhead transcription factor FKHR-L1, which suggests a role of this gene in neuronal and lymphocyte apoptosis. Transgenic studies of the mouse counterpart suggested that this gene functions as an essential initiator of apoptosis in thymocyte-negative selection. Several alternatively spliced transcript variants of this gene have been identified. [provided by RefSeq, Jun 2013]

Immunogen

A synthetic peptide of human Bim

Gene ID

10018

Swiss Prot

O43521

Synonyms

BAM; BIM; BOD

Reactivity

Human,Mouse,Rat

Application

WB, IHC-P, ICC/IF, FC, IP


Recommended dilutionWB: 1:1000
IHC-P: 1:100
ICC/IF: 1:100-1:200
FC: 1:50-1:100
IP: 1:50**Calculated MW**

22 kDa


Observed MW

22 kDa


Host species	Rabbit
Clonality	Monoclonal
Clonality No.	DGR11947
Isotype	IgG
Purity	Affinity Purification
Conjugation	Un-conjugated
Storage Stability	Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.



Western blot analysis of extracts from 3T3, HeLa cells using db13581 at 1:1000.



Western blot analysis of extracts from PC-12 cells and Rat spleen tissue using db13581 at 1:1000.



Immunofluorescence analysis of Jurkat cells labelling Bim with db13581.

The cells were fixed with cold 100% methanol (10min, 4°C) and blocked in 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween 20 for 1h. The cells were then incubate with db13581 (1:100) at room temprature for 1h, followed by a further incubation at room temprature for 45min with Goat Anti Rabbit IgG (H+L)-AF488 (db10005, shown in green). Nuclear DNA was labeled in blue with DAPI.

Control: Secondary antibody only.