

Recombinant

DGRmAb®

**MEK1 (DGR12895) Rabbit mAb**

db13983

Package : 10µL 20µL 50µL 100µL

**Product Name** : MEK1 (DGR12895) Rabbit mAb**Cat.No.:** db13983**Synonyms** : CFC3; MEK1; MKK1; MAPKK1; PRKMK1**Application** : WB, IHC-P, ICC/IF, IP**Reactivity** : Human**Host species** : Rabbit**Background**

The protein encoded by this gene is a member of the dual specificity protein kinase family, which acts as a mitogen-activated protein (MAP) kinase kinase. MAP kinases, also known as extracellular signal-regulated kinases (ERKs), act as an integration point for multiple biochemical signals. This protein kinase lies upstream of MAP kinases and stimulates the enzymatic activity of MAP kinases upon wide variety of extra- and intracellular signals. As an essential component of MAP kinase signal transduction pathway, this kinase is involved in many cellular processes such as proliferation, differentiation, transcription regulation and development. [provided by RefSeq, Jul 2008]

**Immunogen**

A synthetic peptide of human MEK1

**Gene ID**

5604

**Swiss Prot**

Q02750

**Synonyms**

CFC3; MEK1; MKK1; MAPKK1; PRKMK1

**Reactivity**

Human

**Application**

WB, IHC-P, ICC/IF, IP

**Recommended dilution**

WB: 1:1000

IHC-P: 1:50-1:100

ICC/IF: 1:50-1:100

IP: 1:20-1:50

**Calculated MW**

43 kDa

**Observed MW**

43 kDa


**Host species**

Rabbit


**Clonality**

Monoclonal

<b>Clonality No.</b>	DGR12895
<b>Isotype</b>	IgG
<b>Purity</b>	Affinity Purification
<b>Conjugation</b>	Un-conjugated
<b>Storage Stability</b>	Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.



Immunofluorescent analysis of HeLa cells using db13983 antibody (green), and DAPI (blue).



Immunofluorescence analysis of HeLa cells labelling MEK1 with db13983.

The cells were fixed with cold 100% methanol (10min, 4°C) and blocked in 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween 20 for 1h. The cells were then incubate with db13983 (1:50) at room temprature for 1h, followed by a further incubation at room temperature for 45min with Goat Anti Rabbit IgG (H+L)-AF488 (db10005, shown in green). Nuclear DNA was labeled in blue with DAPI.

Control: Secondary antibody only.