

DGRmAb[®]

Recombinant

Cyclin E1 (DGR13531) Rabbit mAb

db14262

Package : 10µL 20µL 50µL 100µL


Product Name : Cyclin E1 (DGR13531) Rabbit mAb Cat.No.: db14262 Synonyms : CCNE; pCCNE1 Application : WB, ICC/IF, FC Reactivity : Human Host species : Rabbit

Background

Background	The protein encoded by this gene belongs to the highly conserved cyclin family, whose members
	are characterized by a dramatic periodicity in protein abundance through the cell cycle. Cyclins
	function as regulators of CDK kinases. Different cyclins exhibit distinct expression and degradation
	patterns which contribute to the temporal coordination of each mitotic event. This cyclin forms a
	complex with and functions as a regulatory subunit of CDK2, whose activity is required for cell cycle
	G1/S transition. This protein accumulates at the G1-S phase boundary and is degraded as cells
	progress through S phase. Overexpression of this gene has been observed in many tumors, which
	results in chromosome instability, and thus may contribute to tumorigenesis. This protein was found
	to associate with, and be involved in, the phosphorylation of NPAT protein (nuclear protein mapped
	to the ATM locus), which participates in cell-cycle regulated histone gene expression and plays a
	critical role in promoting cell-cycle progression in the absence of pRB. [provided by RefSeq, Apr
	2016]
Immunogen	A synthetic peptide of human Cyclin E1
Gene ID	898
Gene ID Swiss Prot	898 P24864
Swiss Prot	P24864
Swiss Prot Synonyms	P24864 CCNE; pCCNE1
Swiss Prot Synonyms Reactivity	P24864 CCNE; pCCNE1 Human
Swiss Prot Synonyms Reactivity Application	P24864 CCNE; pCCNE1 Human WB, ICC/IF, FC
Swiss Prot Synonyms Reactivity Application	P24864 CCNE; pCCNE1 Human WB, ICC/IF, FC WB: 1:1000-1:5000
Swiss Prot Synonyms Reactivity Application	P24864 CCNE; pCCNE1 Human WB, ICC/IF, FC WB: 1:1000-1:5000 ICC/IF: 1:500-1:1000

dvagbvo 戴格生物

Host species	Rabbit
Clonality	Monoclonal
Clonality No.	DGR13531
lsotype	lgG
Purity	Affinity Purification
Conjugation	Un-conjugated
Storage Stability	Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.
456h 488	Western blot detection of Cyclin E1 in K562, Hela cell lysates using Cyclin E1 antibody(1:1000



diluted).