

Recombinant

DGRmAb®

**Cyclin E1 (DGR13531) Rabbit mAb**

db14262

Package : 10µL 20µL 50µL 100µL

**Product Name** : Cyclin E1 (DGR13531) Rabbit mAb**Cat.No.:** db14262**Synonyms** : CCNE; pCCNE1**Application** : WB, ICC/IF, FC**Reactivity** : Human**Host species** : Rabbit**Background**

The protein encoded by this gene belongs to the highly conserved cyclin family, whose members are characterized by a dramatic periodicity in protein abundance through the cell cycle. Cyclins function as regulators of CDK kinases. Different cyclins exhibit distinct expression and degradation patterns which contribute to the temporal coordination of each mitotic event. This cyclin forms a complex with and functions as a regulatory subunit of CDK2, whose activity is required for cell cycle G1/S transition. This protein accumulates at the G1-S phase boundary and is degraded as cells progress through S phase. Overexpression of this gene has been observed in many tumors, which results in chromosome instability, and thus may contribute to tumorigenesis. This protein was found to associate with, and be involved in, the phosphorylation of NPAT protein (nuclear protein mapped to the ATM locus), which participates in cell-cycle regulated histone gene expression and plays a critical role in promoting cell-cycle progression in the absence of pRB. [provided by RefSeq, Apr 2016]

**Immunogen**

A synthetic peptide of human Cyclin E1

**Gene ID**

898

**Swiss Prot**

P24864

**Synonyms**

CCNE; pCCNE1

**Reactivity**

Human

**Application**

WB, ICC/IF, FC

**Recommended dilution**

WB: 1:1000-1:5000

ICC/IF: 1:500-1:1000

FC: 1:100

**Calculated MW**



47 kDa

**Observed MW**

47 kDa

Host species	Rabbit
Clonality	Monoclonal
Clonality No.	DGR13531
Isotype	IgG
Purity	Affinity Purification
Conjugation	Un-conjugated
Storage Stability	Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.

Western blot analysis of extracts from K562, HeLa cells using db14262 at 1:1000.



Immunofluorescence analysis of HeLa cells labelling Cyclin E1 with db14262.

The cells were fixed with 4% PFA (10min, RT) followed by treatment with 0.1% Triton X-100 (10min, RT), and blocked in 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween 20 for 1h. The cells were then incubate with db14262 (1:500) at room temprature for 1h, followed by a further incubation at room temperature for 45min with Goat Anti Rabbit IgG (H+L)-AF488 (db10005, shown in green). Nuclear DNA was labeled in blue with DAPI.

Control: Secondary antibody only.