

Recombinant

DGRmAb®

ICAD (DGR13646) Rabbit mAb

db13371

Package : 10µL 20µL 50µL 100µL

Product Name : ICAD (DGR13646) Rabbit mAb**Cat.No.:** db13371**Synonyms** : DFF1; ICAD; DFF-45**Application** : WB**Reactivity** : Human,Mouse,Rat**Host species** : Rabbit**Background**

Apoptosis is a cell death process that removes toxic and/or useless cells during mammalian development. The apoptotic process is accompanied by shrinkage and fragmentation of the cells and nuclei and degradation of the chromosomal DNA into nucleosomal units. DNA fragmentation factor (DFF) is a heterodimeric protein of 40-kD (DFFB) and 45-kD (DFFA) subunits. DFFA is the substrate for caspase-3 and triggers DNA fragmentation during apoptosis. DFF becomes activated when DFFA is cleaved by caspase-3. The cleaved fragments of DFFA dissociate from DFFB, the active component of DFF. DFFB has been found to trigger both DNA fragmentation and chromatin condensation during apoptosis. Two alternatively spliced transcript variants encoding distinct isoforms have been found for this gene. [provided by RefSeq, Jul 2008]

Immunogen

A synthetic peptide of human ICAD

Gene ID

1676

Swiss Prot

O00273

Synonyms

DFF1; ICAD; DFF-45

Reactivity

Human,Mouse,Rat

Application

WB

Recommended dilution

WB: 1:2000-1:20000

Calculated MW

37 kDa

Observed MW

45 kDa

Host species

Rabbit


Clonality

Monoclonal

Clonality No.

DGR13646

Isotype	IgG
Purity	Affinity Purification
Conjugation	Un-conjugated
Storage Stability	Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.



Western blot analysis of extracts from MCF-7, C6 cells and Rat brain tissue using db13371 at 1:1000.