



Recombinant

DGRmAb[®]

Cyclin H (DGR13708) Rabbit mAb

db14756 Package : 10μL 20μL 50μL 100μL

Product Name: Cyclin H (DGR13708) Rabbit mAb

Cat.No.: db14756

Synonyms: CAK; p34; p37; CycH

Application: WB

Reactivity: Human, Mouse, Rat

Host species: Rabbit

Background The protein encoded by this gene belongs to the highly conserved cyclin family, whose members

are characterized by a dramatic periodicity in protein abundance through the cell cycle. Cyclins function as regulators of CDK kinases. Different cyclins exhibit distinct expression and degradation patterns which contribute to the temporal coordination of each mitotic event. This cyclin forms a complex with CDK7 kinase and ring finger protein MAT1. The kinase complex is able to phosphorylate CDK2 and CDC2 kinases, thus functions as a CDK-activating kinase (CAK). This cyclin and its kinase partner are components of TFIIH, as well as RNA polymerase II protein complexes. They participate in two different transcriptional regulation processes, suggesting an important link between basal transcription control and the cell cycle machinery. A pseudogene of

this gene is found on chromosome 4. Alternate splicing results in multiple transcript variants.

[provided by RefSeq, Nov 2010]

Immunogen A synthetic peptide of human Cyclin H

Gene ID 902

Swiss Prot P51946

Synonyms CAK; p34; p37; CycH

Reactivity Human, Mouse, Rat

Application WB

Recommended dilution WB: 1:1000-1:5000

Calculated MW 38 kDa

Observed MW 36 kDa

Host species Rabbit

Clonality Monoclonal





Clonality No. DGR13708


Isotype IgG

Purity Affinity Purification

Conjugation Un-conjugated

Storage Stability Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium

azide and 0.05% BSA. Stable for 12 months from date of receipt.



Western blot detection of Cyclin H in Jurkat cell lysates using Cyclin H antibody(1:1000 diluted).