

Recombinant DGRmAb<sup>®</sup>

HDAC2 (DGR14124) Rabbit mAb

db11911

Package : 10µL 20µL 50µL 100µL


Product Name : HDAC2 (DGR14124) Rabbit mAb Cat.No.: db11911 Synonyms : HD2; RPD3; YAF1 Application : WB, IHC-P, ICC/IF, FC, IP Reactivity : Human,Mouse,Rat Host species : Rabbit

| Background           | This gene product belongs to the histone deacetylase family. Histone deacetylases act via the formation of large multiprotein complexes, and are responsible for the deacetylation of lysine residues at the N-terminal regions of core histones (H2A, H2B, H3 and H4). This protein forms transcriptional repressor complexes by associating with many different proteins, including YY1, a mammalian zinc-finger transcription factor. Thus, it plays an important role in transcriptional regulation, cell cycle progression and developmental events. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Apr 2010] |
|----------------------|--|
| Immunogen            | Recombinant protein of human HDAC2   |
| Gene ID              | 3066   |
| Swiss Prot           | Q92769   |
| Synonyms             | HD2; RPD3; YAF1  |
| Reactivity           | Human,Mouse,Rat  |
| Application          | WB, IHC-P, ICC/IF, FC, IP  |
| Recommended dilution | WB: 1:1000<br>IHC-P: 1:500-1:2000<br>ICC/IF: 1:500-1:1000<br>FC: 1:200-1:500<br>IP: 1:20-1:50  |
| Calculated MW        | 55 kDa   |
| Observed MW          | 60 kDa   |
| Host species         | Rabbit   |
| Clonality            | Monoclonal   |


## dvagbvo 戴格生物

| Clonality No.     | DGR14124  |
|-------------------|---|
| lsotype           | lgG   |
| Purity            | Affinity Purification   |
| Conjugation       | Un-conjugated   |
| Storage Stability | Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt. |







Western blot detection of HDAC2 in K562,C6,3T3 cell lysates using HDAC2 antibody(1:1000 diluted).



Western blot analysis of extracts from C2C12 cells and Rat muscle tissue using db11911 at 1:1000.



Immunohistochemical analysis of paraffin-embedded human colon cancer using db11911 antibody.



Immunofluorescent analysis of HeLa cells using db11911 antibody (green), and DAPI (blue).

