

Recombinant

DGRmAb®

BRG1 (DGR14531) Rabbit mAb

db11111

Package : 10µL 20µL 50µL 100µL

Product Name : BRG1 (DGR14531) Rabbit mAb**Cat.No.:** db11111**Synonyms** : Brg1; SW1/SNF; HP1-BP72; SNF2beta; b2b692Clo; b2b508.1Clo**Application** : WB, IHC-P, ICC/IF, FC, IP**Reactivity** : Human,Mouse,Rat**Host species** : Rabbit**Background**

Transcriptional coactivator cooperating with nuclear hormone receptors to potentiate transcriptional activation. Component of the CREST-BRG1 complex, a multiprotein complex that regulates promoter activation by orchestrating a calcium-dependent release of a repressor complex and a recruitment of an activator complex. In resting neurons, transcription of the c-FOS promoter is inhibited by BRG1-dependent recruitment of a phospho-RB1-HDAC repressor complex. Upon calcium influx, RB1 is dephosphorylated by calcineurin, which leads to release of the repressor complex. At the same time, there is increased recruitment of CREBBP to the promoter by a CREST-dependent mechanism, which leads to transcriptional activation. The CREST-BRG1 complex also binds to the NR2B promoter, and activity-dependent induction of NR2B expression involves a release of HDAC1 and recruitment of CREBBP (By similarity).

Immunogen

A synthetic peptide of mouse BRG1

Gene ID

20586

Swiss Prot

Q3TKT4

Synonyms

Brg1; SW1/SNF; HP1-BP72; SNF2beta; b2b692Clo; b2b508.1Clo

Reactivity

Human,Mouse,Rat

Application

WB, IHC-P, ICC/IF, FC, IP

Recommended dilution

WB: 1:2000-1:20000

IHC-P: 1:100-1:200

ICC/IF: 1:200-1:500

FC: 1:100-1:200

IP: 1:10-1:100


Calculated MW

185 kDa


Observed MW

220 kDa


Host species	Rabbit
Clonality	Monoclonal
Clonality No.	DGR14531
Isotype	IgG
Purity	Affinity Purification
Conjugation	Un-conjugated
Storage Stability	Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.



Western blot analysis of extracts from 293T, HeLa, A549 cells using db11111 at 1:1000.



Western blot analysis of extracts from 3T3, C6 cells using db11111 at 1:1000.



Immunofluorescence analysis of HeLa cells labelling BRG1 with db11111.

The cells were fixed with 4% PFA (10min, RT) followed by treatment with 0.1% Triton X-100 (10min, RT), and blocked in 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween 20 for 1h. The cells were then incubate with db11111 (1:200) at room temprature for 1h, followed by a further incubation at room temperature for 45min with Goat Anti Rabbit IgG (H+L)-AF488 (db10005, shown in green). Nuclear DNA was labeled in blue with DAPI.

Control: Secondary antibody only.