Recombinant DGRmAb®


TRAF6 (DGR14750) Rabbit mAb

db12997 **Package : 10μL 20μL 50μL 100μL**


Product Name : TRAF6 (DGR14750) Rabbit mAb
Cat.No.: db12997
Synonyms : RNF85; MGC:3310
Application : WB, IHC-P, ICC/IF
Reactivity : Human,Mouse,Rat
Host species : Rabbit

Background	The protein encoded by this gene is a member of the TNF receptor associated factor (TRAF) protein family. TRAF proteins are associated with, and mediate signal transduction from, members of the TNF receptor superfamily. This protein mediates signaling from members of the TNF receptor superfamily as well as the Toll/IL-1 family. Signals from receptors such as CD40, TNFSF11/RANCE and IL-1 have been shown to be mediated by this protein. This protein also interacts with various protein kinases including IRAK1/IRAK, SRC and PKCzeta, which provides a link between distinct signaling pathways. This protein functions as a signal transducer in the NF-kappaB pathway that activates IkappaB kinase (IKK) in response to proinflammatory cytokines. The interaction of this protein with UBE2N/UBC13, and UBE2V1/UEV1A, which are ubiquitin conjugating enzymes catalyzing the formation of polyubiquitin chains, has been found to be required for IKK activation by this protein. This protein also interacts with the transforming growth factor (TGF) beta receptor complex and is required for Smad-independent activation of the JNK and p38 kinases. This protein has an amino terminal RING domain which is followed by four zinc-finger motifs, a central coiled-coil region and a highly conserved carboxyl terminal domain, known as the TRAF-C domain. Two alternatively spliced transcript variants, encoding an identical protein, have been reported. [provided by RefSeq, Feb 2012]
Immunogen	A synthetic peptide of human TRAF6
Gene ID	7189, 22034, 311245
Swiss Prot	Q9Y4K3, P70196, B5DF45
Synonyms	RNF85; MGC:3310
Reactivity	Human,Mouse,Rat
Application	WB, IHC-P, ICC/IF
Recommended dilution	WB: 1:1000-1:5000 IHC-P: 1:50

	ICC/IF: 1:50
Calculated MW	60 kDa
Observed MW	60 kDa
Host species	Rabbit
Clonality	Monoclonal
Clonality No.	DGR14750
Isotype	IgG
Purity	Affinity Purification
Conjugation	Un-conjugated
Storage Stability	Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.



Western blot analysis of extracts from K562, C6, HeLa cells and Rat brain tissue using db12997 at 1:1000.



Immunofluorescence analysis of HeLa cells labelling TRAF6 with db12997.

The cells were fixed with 4% PFA (10min, RT) followed by treatment with 0.1% Triton X-100 (10min, RT), and blocked in 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween 20 for 1h. The cells were then incubate with db12997 (1:50) at room temprature for 1h, followed by a further incubation at room temperature for 45min with Goat Anti Rabbit IgG (H+L)-AF488 (db10005, shown in green). Nuclear DNA was labeled in blue with DAPI.

Control: Secondary antibody only.