Recombinant DGRmAb®


p38 (DGR14777) Rabbit mAb

db14381 **Package : 10μL 20μL 50μL 100μL**


Product Name : p38 (DGR14777) Rabbit mAb
Cat.No.: db14381
Synonyms : RK; p38; CSBP; EXIP; Mxi2; CSBP1; CSBP2; CSPB1; PRKM14; PRKM15; SAPK2A; p38ALPHA
Application : WB, IHC, ICC/IF
Reactivity : Human, Mouse, Rat
Host species : Rabbit

Background	The protein encoded by this gene is a member of the MAP kinase family. MAP kinases act as an integration point for multiple biochemical signals, and are involved in a wide variety of cellular processes such as proliferation, differentiation, transcription regulation and development. This kinase is activated by various environmental stresses and proinflammatory cytokines. The activation requires its phosphorylation by MAP kinase kinases (MKKs), or its autophosphorylation triggered by the interaction of MAP3K7IP1/TAB1 protein with this kinase. The substrates of this kinase include transcription regulator ATF2, MEF2C, and MAX, cell cycle regulator CDC25B, and tumor suppressor p53, which suggest the roles of this kinase in stress related transcription and cell cycle regulation, as well as in genotoxic stress response. Four alternatively spliced transcript variants of this gene encoding distinct isoforms have been reported. [provided by RefSeq, Jul 2008]
Immunogen	A synthetic peptide of human p38
Gene ID	1432
Swiss Prot	Q16539
Synonyms	RK; p38; CSBP; EXIP; Mxi2; CSBP1; CSBP2; CSPB1; PRKM14; PRKM15; SAPK2A; p38ALPHA
Reactivity	Human, Mouse, Rat
Application	WB, IHC, ICC/IF
Recommended dilution	WB: 1:1000-1:5000 IHC: 1:100-1:200 ICC/IF: 1:100-1:200
Calculated MW	41 kDa
Observed MW	41 kDa

Host species	Rabbit
Clonality	Monoclonal
Clonality No.	DGR14777
Isotype	IgG
Purity	Affinity Purification
Conjugation	Un-conjugated
Storage Stability	Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.



Western blot analysis of extracts from Jurkat, C6, CHO-K1, HeLa cells using db14381 at 1:1000.



Immunofluorescence analysis of HeLa cells labelling p38 with db14381.

The cells were fixed with 4% PFA (10min, RT) followed by treatment with 0.1% Triton X-100 (10min, RT), and blocked in 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween 20 for 1h. The cells were then incubate with db14381 (1:100) at room temprature for 1h, followed by a further incubation at room temperature for 45min with Goat Anti Rabbit IgG (H+L)-AF488 (db10005, shown in green). Nuclear DNA was labeled in blue with DAPI.

Control: Secondary antibody only.