

Recombinant

DGRmAb®

p38 (DGR14777) Rabbit mAb

db14381

Package : 10µL 20µL 50µL 100µL

Product Name : p38 (DGR14777) Rabbit mAb**Cat.No.:** db14381**Synonyms :** RK; p38; CSBP; EXIP; Mxi2; CSBP1; CSBP2; CSPB1; PRKM14; PRKM15; SAPK2A; p38ALPHA**Application :** WB, IHC, ICC/IF**Reactivity :** Human, Mouse, Rat**Host species :** Rabbit**Background**

The protein encoded by this gene is a member of the MAP kinase family. MAP kinases act as an integration point for multiple biochemical signals, and are involved in a wide variety of cellular processes such as proliferation, differentiation, transcription regulation and development. This kinase is activated by various environmental stresses and proinflammatory cytokines. The activation requires its phosphorylation by MAP kinase kinases (MKKs), or its autophosphorylation triggered by the interaction of MAP3K7IP1/TAB1 protein with this kinase. The substrates of this kinase include transcription regulator ATF2, MEF2C, and MAX, cell cycle regulator CDC25B, and tumor suppressor p53, which suggest the roles of this kinase in stress related transcription and cell cycle regulation, as well as in genotoxic stress response. Four alternatively spliced transcript variants of this gene encoding distinct isoforms have been reported. [provided by RefSeq, Jul 2008]

Immunogen

A synthetic peptide of human p38

Gene ID

1432

Swiss Prot

Q16539

Synonyms

RK; p38; CSBP; EXIP; Mxi2; CSBP1; CSBP2; CSPB1; PRKM14; PRKM15; SAPK2A; p38ALPHA

Reactivity

Human, Mouse, Rat

Application

WB, IHC, ICC/IF

Recommended dilution

WB: 1:1000-1:5000

IHC: 1:100-1:200

ICC/IF: 1:100-1:200


Calculated MW

41 kDa

Observed MW

41 kDa

Host species	Rabbit
Clonality	Monoclonal
Clonality No.	DGR14777
Isotype	IgG
Purity	Affinity Purification
Conjugation	Un-conjugated
Storage Stability	Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.



Western blot detection of p38 in Jurkat,C6,CHO-K1,Hela cell lysates using p38 antibody(1:500 diluted).