

Recombinant

DGRmAb®

Caspase-3 p12 (DGR16337) Rabbit mAb

db11371

Package : 10µL 20µL 50µL 100µL

Product Name : Caspase-3 p12 (DGR16337) Rabbit mAb**Cat.No.:** db11371**Synonyms** : CPP32; SCA-1; CPP32B**Application** : WB, IHC-P, ICC/IF, FC**Reactivity** : Human,Mouse,Rat**Host species** : Rabbit**Background**

The protein encoded by this gene is a cysteine-aspartic acid protease that plays a central role in the execution-phase of cell apoptosis. The encoded protein cleaves and inactivates poly(ADP-ribose) polymerase while it cleaves and activates sterol regulatory element binding proteins as well as caspases 6, 7, and 9. This protein itself is processed by caspases 8, 9, and 10. It is the predominant caspase involved in the cleavage of amyloid-beta 4A precursor protein, which is associated with neuronal death in Alzheimer's disease. [provided by RefSeq, Aug 2017]

Immunogen

Recombinant protein of human Caspase-3 p12

Gene ID

836

Swiss Prot

P42574

Synonyms

CPP32; SCA-1; CPP32B

Reactivity

Human,Mouse,Rat

Application

WB, IHC-P, ICC/IF, FC

Recommended dilutionWB: 1:1000
IHC-P: 1:200-1:2000
ICC/IF: 1:100
FC: 1:100-1:200**Calculated MW**

32 kDa

Observed MW

32,12 kDa

Host species

Rabbit


Clonality

Monoclonal


Clonality No.

DGR16337

Isotype	IgG
Purity	Affinity Purification
Conjugation	Un-conjugated
Storage Stability	Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.



Western blot analysis of extracts from K562,3T3,HeLa cells using db11371 at 1:1000.



Immunofluorescence analysis of HeLa cells labelling Caspase-3 p12 with db11371.

The cells were fixed with cold 100% methanol (10min, 4°C) and blocked in 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween 20 for 1h. The cells were then incubate with db11371 (1:100) at room temprature for 1h, followed by a further incubation at room temperature for 45min with Goat Anti Rabbit IgG (H+L)-AF488 (db10005, shown in green). Nuclear DNA was labeled in blue with DAPI.

Control: Secondary antibody only.