

Recombinant

DGRmAb®

## Hexokinase 1 (DGR16439) Rabbit mAb

db11509

Package : 10µL 20µL 50µL 100µL

**Product Name** : Hexokinase 1 (DGR16439) Rabbit mAb

**Cat.No.:** db11509

**Synonyms** : HK; HKD; HKI; HXK1; RP79; HMSNR; HK1-ta; HK1-tb; HK1-tc; hexokinase

**Application** : WB, IHC-P, ICC/IF, FC

**Reactivity** : Human,Mouse,Rat

**Host species** : Rabbit

### Background

Hexokinases phosphorylate glucose to produce glucose-6-phosphate, the first step in most glucose metabolism pathways. This gene encodes a ubiquitous form of hexokinase which localizes to the outer membrane of mitochondria. Mutations in this gene have been associated with hemolytic anemia due to hexokinase deficiency. Alternative splicing of this gene results in several transcript variants which encode different isoforms, some of which are tissue-specific. [provided by RefSeq, Apr 2016]

### Immunogen

A synthetic peptide of human Hexokinase 1

### Gene ID

3098

### Swiss Prot

P19367

### Synonyms

HK; HKD; HKI; HXK1; RP79; HMSNR; HK1-ta; HK1-tb; HK1-tc; hexokinase

### Reactivity

Human,Mouse,Rat

### Application

WB, IHC-P, ICC/IF, FC

### Recommended dilution

WB: 1:1000-1:5000

IHC-P: 1:50-1:100

ICC/IF: 1:50

FC: 1:20

### Calculated MW

103 kDa

### Observed MW

103 kDa

### Host species

Rabbit


### Clonality

Monoclonal


### Clonality No.

DGR16439


<b>Isotype</b>	IgG
<b>Purity</b>	Affinity Purification
<b>Conjugation</b>	Un-conjugated
<b>Storage Stability</b>	Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.




Western blot analysis of extracts from K562, 3T3 cells and Rat brain tissue using db11509 at 1:1000.





Western blot analysis of extracts from HeLa cells and Mouse heart tissue using db11509 at 1:2000.



Immunohistochemical analysis of paraffin-embedded human tonsil using db11509 antibody.



Immunofluorescent analysis of K562 cells using db11509 antibody (green), and DAPI (blue).



Immunofluorescence analysis of HeLa cells labelling Hexokinase 1 with db11509.

The cells were fixed with 4% PFA (10min, RT) followed by treatment with 0.1% Triton X-100 (10min, RT), and blocked in 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween 20 for 1h. The cells were then incubate with db11509 (1:50) at room temprature for 1h, followed by a further incubation at room temperature for 45min with Goat Anti Rabbit IgG (H+L)-AF488 (db10005, shown in green). Nuclear DNA was labeled in blue with DAPI.

Control: Secondary antibody only.