

Recombinant

DGRmAb®

## GAPDH (DGR18773) Rabbit mAb

db14510

Package : 10µL 20µL 100µL 500µL 1mL

**Product Name :** GAPDH (DGR18773) Rabbit mAb

**Cat.No.:** db14510

**Synonyms :** G3PD; GAPD; HEL-S-162eP

**Application :** WB, IHC-P, ICC/IF, FC

**Reactivity :** Human,Mouse,Rat,Monkey,Cow, Dog,Chicken

**Host species :** Rabbit

### Background

This gene encodes a member of the glyceraldehyde-3-phosphate dehydrogenase protein family. The encoded protein has been identified as a moonlighting protein based on its ability to perform mechanistically distinct functions. The product of this gene catalyzes an important energy-yielding step in carbohydrate metabolism, the reversible oxidative phosphorylation of glyceraldehyde-3-phosphate in the presence of inorganic phosphate and nicotinamide adenine dinucleotide (NAD). The encoded protein has additionally been identified to have uracil DNA glycosylase activity in the nucleus. Also, this protein contains a peptide that has antimicrobial activity against *E. coli*, *P. aeruginosa*, and *C. albicans*. Studies of a similar protein in mouse have assigned a variety of additional functions including nitrosylation of nuclear proteins, the regulation of mRNA stability, and acting as a transferrin receptor on the cell surface of macrophage. Many pseudogenes similar to this locus are present in the human genome. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Nov 2014]

### Immunogen

Recombinant protein of human GAPDH

### Gene ID

2597

### Swiss Prot

P04406

### Synonyms

G3PD; GAPD; HEL-S-162eP

### Reactivity

Human,Mouse,Rat,Monkey,Cow, Dog,Chicken

### Application

WB, IHC-P, ICC/IF, FC


### Recommended dilution

WB: 1:1000-1:5000  
 IHC-P: 1:200-1:500  
 ICC/IF: 1:100-1:200  
 FC: 1:100-1:200


### Calculated MW

36 kDa

Observed MW	36 kDa
Host species	Rabbit
Clonality	Monoclonal
Clonality No.	DGR18773
Isotype	IgG
Purity	Affinity Purification
Conjugation	Un-conjugated
Storage Stability	Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.



Western blot analysis of extracts from HeLa, CHO-K1, C6, Jurkat cells and Rat brain tissue using db14510 at 1:1000.



Immunofluorescence analysis of HeLa cells labelling GAPDH with db14510.

The cells were fixed with cold 100% methanol (10min, 4°C) and blocked in 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween 20 for 1h. The cells were then incubated with db14510 (1:100) at room temperature for 1h, followed by a further incubation at room temperature for 45min with Goat Anti Rabbit IgG (H+L)-AF488 (db10005, shown in green). Nuclear DNA was labeled in blue with DAPI.

Control: Secondary antibody only.