

Recombinant DGRmAb[®]

Phospho-ERK1/2 (Thr202/Thr185) (DGR12177) Rabbit mAb

db14364

Package : 10µL 20µL 50µL 100µL

Product Name : Phospho-ERK1/2 (Thr202/Thr185) (DGR12177) Rabbit mAb
Cat.No.: db14364
Synonyms : ERK1; ERT2; ERK-1; PRKM3; P44ERK1; P44MAPK; HS44KDAP; HUMKER1A; p44-ERK1; p44-MAPK
Application : WB, IHC-P, ICC/IF, IP
Reactivity : Human,Mouse,Rat

Host species : Rabbit

Background	The protein encoded by this gene is a member of the MAP kinase family. MAP kinases, also
	known as extracellular signal-regulated kinases (ERKs), act in a signaling cascade that regulates
	various cellular processes such as proliferation, differentiation, and cell cycle progression in
	response to a variety of extracellular signals. This kinase is activated by upstream kinases,
	resulting in its translocation to the nucleus where it phosphorylates nuclear targets. Alternatively
	spliced transcript variants encoding different protein isoforms have been described. [provided by
	RefSeq, Jul 2008]
Immunogen	A synthetic phosphopeptide corresponding to residues surrounding Thr202 of human ERK1
Gene ID	5595
Swiss Prot	P27361
Synonyms	ERK1; ERT2; ERK-1; PRKM3; P44ERK1; P44MAPK; HS44KDAP; HUMKER1A; p44-ERK1;
	p44-MAPK
Reactivity	Human,Mouse,Rat
Application	WB, IHC-P, ICC/IF, IP
Recommended dilution	WB: 1:1000
	IHC-P: 1:200-1:500
	ICC/IF: 1:100
	IP: 1:20-1:50
Calculated MW	43,41 kDa
Observed MW	44,42 kDa
Host species	Rabbit
Clonality	Monoclonal

dvagbvo 戴格生物

Clonality No.	DGR12177
lsotype	lgG
Purity	Affinity Purification
Conjugation	Un-conjugated
Storage Stability	Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.
180- 130- 130- 95- 72- 55- 43- 24- 26- 17-	Western blot detection of Phospho-p44/42 MAPK (ERK1/2) (Thr202) in C6,Hela cell lysates using Phospho-p44/42 MAPK (ERK1/2) (Thr202) antibody(1:1000 diluted).