

Recombinant

DGRmAb®

## Caspase-7 (DGR33388) Rabbit mAb

db14082

Package : 10µL 20µL 50µL 100µL

**Product Name** : Caspase-7 (DGR33388) Rabbit mAb**Cat.No.:** db14082**Synonyms** : MCH3; CMH-1; LICE2; CASP-7; ICE-LAP3**Application** : WB, IHC-P**Reactivity** : Human**Host species** : Rabbit**Background**

This gene encodes a member of the cysteine-aspartic acid protease (caspase) family. Sequential activation of caspases plays a central role in the execution-phase of cell apoptosis. Caspases exist as inactive proenzymes which undergo proteolytic processing at conserved aspartic residues to produce two subunits, large and small, that dimerize to form the active enzyme. The precursor of the encoded protein is cleaved by caspase 3 and 10, is activated upon cell death stimuli and induces apoptosis. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene. [provided by RefSeq, May 2012]

**Immunogen**

A synthetic peptide of human pro Caspase-7

**Gene ID**

840

**Swiss Prot**

P55210

**Synonyms**

MCH3; CMH-1; LICE2; CASP-7; ICE-LAP3

**Reactivity**

Human

**Application**

WB, IHC-P

**Recommended dilution**

WB: 1:1000

IHC-P: 1:50

**Calculated MW**

34 kDa

**Observed MW**

34 kDa

**Host species**

Rabbit

**Clonality**

Monoclonal


**Clonality No.**

DGR33388


**Isotype**

IgG

Purity	Affinity Purification
Conjugation	Un-conjugated
Storage Stability	Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.



Western blot detection of pro Caspase-7 in A549,MCF-7,293T cell lysates using pro Caspase-7 antibody(1:1000 diluted).



Immunohistochemical analysis of paraffin-embedded human Cholangiocarcinoma using db14082 antibody.