Recombinant DGRmAb®

Histone H2A.X (DGR20837) Rabbit mAb

db13705 **Package : 10μL 20μL 50μL 100μL**

Product Name : Histone H2A.X (DGR20837) Rabbit mAb
Cat.No.: db13705
Synonyms : H2A.X; H2A/X; H2AFX
Application : WB, IHC-P, ICC/IF, FC, IP
Reactivity : Human
Host species : Rabbit

Background	Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene encodes a replication-independent histone that is a member of the histone H2A family, and generates two transcripts through the use of the conserved stem-loop termination motif, and the polyA addition motif. [provided by RefSeq, Oct 2015]
Immunogen	A synthetic peptide of human Histone H2A.X
Gene ID	3014
Swiss Prot	P16104
Synonyms	H2A.X; H2A/X; H2AFX
Reactivity	Human
Application	WB, IHC-P, ICC/IF, FC, IP
Recommended dilution	WB: 1:1000 IHC-P: 1:100-1:200 ICC/IF: 1:100 FC: 1:20-1:50 IP: 1:20-1:50
Calculated MW	15 kDa
Observed MW	15 kDa
Host species	Rabbit

Clonality	Monoclonal
Clonality No.	DGR20837
Isotype	IgG
Purity	Affinity Purification
Conjugation	Un-conjugated
Storage Stability	Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.



Western blot detection of Histone H2A.X in Hela,Ramos using Histone H2A.X antibody(1:1000 diluted)

Western blot analysis of extracts from Jurkat cells using db13705 at 1:1000.





Immunohistochemical analysis of paraffin-embedded human lung cancer tissue using db13705 antibody.



Immunofluorescent analysis of HeLa cells using db13705 antibody (green), and DAPI (blue).