Recombinant DGRmAb®


HDAC9 (DGR21105) Rabbit mAb

db14340 **Package : 10μL 20μL 50μL 100μL**


Product Name : HDAC9 (DGR21105) Rabbit mAb
Cat.No.: db14340
Synonyms : HD7; HD9; HD7b; HDAC; HDRP; MITR; HDAC7; HDAC7B; HDAC9B; HDAC9FL
Application : WB, IHC-P, ICC/IF, FC, IP
Reactivity : Human
Host species : Rabbit

Background	Histones play a critical role in transcriptional regulation, cell cycle progression, and developmental events. Histone acetylation/deacetylation alters chromosome structure and affects transcription factor access to DNA. The protein encoded by this gene has sequence homology to members of the histone deacetylase family. This gene is orthologous to the Xenopus and mouse MITR genes. The MITR protein lacks the histone deacetylase catalytic domain. It represses MEF2 activity through recruitment of multicomponent corepressor complexes that include CtBP and HDACs. This encoded protein may play a role in hematopoiesis. Multiple alternatively spliced transcripts have been described for this gene but the full-length nature of some of them has not been determined. [provided by RefSeq, Jul 2008]
Immunogen	A synthetic peptide of human HDAC9
Gene ID	9734
Swiss Prot	Q9UKV0
Synonyms	HD7; HD9; HD7b; HDAC; HDRP; MITR; HDAC7; HDAC7B; HDAC9B; HDAC9FL
Reactivity	Human
Application	WB, IHC-P, ICC/IF, FC, IP
Recommended dilution	WB: 1:1000-1:5000 IHC-P: 1:200-1:1000 ICC/IF: 1:100-1:200 FC: 1:100 IP: 1:20-1:50
Calculated MW	111 kDa
Observed MW	150 kDa
Host species	Rabbit


Clonality	Monoclonal
Clonality No.	DGR21105
Isotype	IgG
Purity	Affinity Purification
Conjugation	Un-conjugated
Storage Stability	Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.



Western blot analysis of extracts from HeLa, HepG2 cells using db14340 at 1:4000.



Immunohistochemical analysis of paraffin-embedded human tonsil using db14340 antibody.



Immunofluorescence analysis of HeLa cells labelling HDAC9 with db14340.

The cells were fixed with 4% PFA (10min, RT) followed by treatment with 0.1% Triton X-100 (10min, RT), and blocked in 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween 20 for 1h. The cells were then incubate with db14340 (1:100) at room temprature for 1h, followed by a further incubation at room temperature for 45min with Goat Anti Rabbit IgG (H+L)-AF488 (db10005, shown in green). Nuclear DNA was labeled in blue with DAPI.

Control: Secondary antibody only.