

Recombinant

DGRmAb®

Phospho-gamma H2A.X (Ser139) (DGR18728) Rabbit mAb

db13191

Package : 10µL 20µL 50µL 100µL

Product Name : Phospho-gamma H2A.X (Ser139) (DGR18728) Rabbit mAb**Cat.No.:** db13191**Synonyms** : H2AX; H2A.X; H2A/X**Application** : WB, IHC-P, ICC/IF, IP**Reactivity** : Human,Mouse,Rat**Host species** : Rabbit**Background**

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene encodes a replication-independent histone that is a member of the histone H2A family, and generates two transcripts through the use of the conserved stem-loop termination motif, and the polyA addition motif. [provided by RefSeq, Oct 2015]

Immunogen

A synthetic phosphopeptide corresponding to residues surrounding Ser139 of human gamma H2A.X

Gene ID

3014

Swiss Prot

P16104

Synonyms

H2AX; H2A.X; H2A/X

Reactivity

Human,Mouse,Rat

Application

WB, IHC-P, ICC/IF, IP

Recommended dilution

WB: 1:1000-1:5000

IHC-P: 1:200-1:1000

ICC/IF: 1:100-1:200

IP: 1:20-1:50

Calculated MW

15 kDa

Observed MW

15 kDa


Host species

Rabbit


Clonality

Monoclonal

Clonality No.	DGR18728
Isotype	IgG
Purity	Affinity Purification
Conjugation	Un-conjugated
Storage Stability	Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.



Western blot analysis of extracts from Jurkat, THP-1, C6, 3T3 cells using db13191 at 1:1000.



Western blot analysis of extracts from HepG2 cells using db13191 at 1:1000.