







MCM2 (DGR20534) Rabbit mAb

db11270 Package : 10μL 20μL 50μL 100μL

Product Name: MCM2 (DGR20534) Rabbit mAb

Cat.No.: db11270

Synonyms: BM28; CCNL1; CDCL1; cdc19; DFNA70; D3S3194; MITOTIN

Application : WB, IHC, ICC/IF, FC **Reactivity :** Human, Mouse, Rat

Host species: Rabbit

Background The protein encoded by this gene is one of the highly conserved mini-chromosome maintenance

proteins (MCM) that are involved in the initiation of eukaryotic genome replication. The hexameric protein complex formed by MCM proteins is a key component of the pre-replication complex (pre_RC) and may be involved in the formation of replication forks and in the recruitment of other DNA replication related proteins. This protein forms a complex with MCM4, 6, and 7, and has been shown to regulate the helicase activity of the complex. This protein is phosphorylated, and thus regulated by, protein kinases CDC2 and CDC7. Multiple alternatively spliced transcript variants

have been found, but the full-length nature of some variants has not been defined. [provided by

RefSeq, Oct 2012]

Immunogen A synthetic peptide of human MCM2

Gene ID 4171

Swiss Prot P49736

Synonyms BM28; CCNL1; CDCL1; cdc19; DFNA70; D3S3194; MITOTIN

Reactivity Human, Mouse, Rat

Application WB, IHC, ICC/IF, FC

Recommended dilution WB: 1:1000-1:5000

IHC: 1:200

ICC/IF: 1:200-1:1000

FC: 1:100-1:200

Calculated MW 102 kDa

Observed MW 125 kDa

Host species Rabbit





Clonality Monoclonal

Clonality No. DGR20534


Isotype IgG

Purity Affinity Purification


Conjugation Un-conjugated

Storage Stability Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium

azide and 0.05% BSA. Stable for 12 months from date of receipt.



Western blot detection of MCM2 in K562,C6,3T3,Hela cell lysates using MCM2 antibody(1:1000 diluted).



Immunofluorescent analysis of HeLa cells using db11270 antibody (green), and DAPI (blue).