

Recombinant

DGRmAb®

## Phospho-Smad2 (Ser250) (DGR31400) Rabbit mAb

db11432

Package : 10µL 20µL 50µL 100µL

**Product Name :** Phospho-Smad2 (Ser250) (DGR31400) Rabbit mAb**Cat.No.:** db11432**Synonyms :** JV18; MADH2; MADR2; JV18-1; hMAD-2; hSMAD2**Application :** WB, FC**Reactivity :** Human,Mouse,Rat**Host species :** Rabbit**Background**

The protein encoded by this gene belongs to the SMAD, a family of proteins similar to the gene products of the Drosophila gene 'mothers against decapentaplegic' (Mad) and the C. elegans gene Sma. SMAD proteins are signal transducers and transcriptional modulators that mediate multiple signaling pathways. This protein mediates the signal of the transforming growth factor (TGF)-beta, and thus regulates multiple cellular processes, such as cell proliferation, apoptosis, and differentiation. This protein is recruited to the TGF-beta receptors through its interaction with the SMAD anchor for receptor activation (SARA) protein. In response to TGF-beta signal, this protein is phosphorylated by the TGF-beta receptors. The phosphorylation induces the dissociation of this protein with SARA and the association with the family member SMAD4. The association with SMAD4 is important for the translocation of this protein into the nucleus, where it binds to target promoters and forms a transcription repressor complex with other cofactors. This protein can also be phosphorylated by activin type 1 receptor kinase, and mediates the signal from the activin. Alternatively spliced transcript variants have been observed for this gene. [provided by RefSeq, May 2012]

**Immunogen**

A synthetic phosphopeptide corresponding to residues surrounding Ser250 of human Smad2

**Gene ID**

4087

**Swiss Prot**

Q15796

**Synonyms**

JV18; MADH2; MADR2; JV18-1; hMAD-2; hSMAD2

**Reactivity**

Human,Mouse,Rat

**Application**

WB, FC

**Recommended dilution**


WB: 1:1000-1:5000

FC: 1:20-1:50


**Calculated MW**

52,48 kDa

Observed MW	58,62 kDa
Host species	Rabbit
Clonality	Monoclonal
Clonality No.	DGR31400
Isotype	IgG
Purity	Affinity Purification
Conjugation	Un-conjugated
Storage Stability	Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.



Western blot detection of Smad2 (phospho-Ser250) in HeLa,CHO-K1,C6 using Smad2 (phospho-Ser250) antibody(1:1000 diluted).



Western blot analysis of extracts from Mouse kidney tissue using db11432 at 1:1000.