

Recombinant

DGRmAb®

DDIT3 (DGR14242) Rabbit mAb

db11608

Package : 10µL 20µL 50µL 100µL

Product Name : DDIT3 (DGR14242) Rabbit mAb**Cat.No.:** db11608**Synonyms** : CHOP; CEBPZ; CHOP10; CHOP-10; GADD153; C/EBPzeta**Application** : WB, IHC-P**Reactivity** : Human,Mouse,Rat**Host species** : Rabbit**Background**

This gene encodes a member of the CCAAT/enhancer-binding protein (C/EBP) family of transcription factors. The protein functions as a dominant-negative inhibitor by forming heterodimers with other C/EBP members, such as C/EBP and LAP (liver activator protein), and preventing their DNA binding activity. The protein is implicated in adipogenesis and erythropoiesis, is activated by endoplasmic reticulum stress, and promotes apoptosis. Fusion of this gene and FUS on chromosome 16 or EWSR1 on chromosome 22 induced by translocation generates chimeric proteins in myxoid liposarcomas or Ewing sarcoma. Multiple alternatively spliced transcript variants encoding two isoforms with different length have been identified. [provided by RefSeq, Aug 2010]

Immunogen

A synthetic peptide of human DDIT3

Gene ID

1649

Swiss Prot

P35638

Synonyms

CHOP; CEBPZ; CHOP10; CHOP-10; GADD153; C/EBPzeta

Reactivity

Human,Mouse,Rat

Application

WB, IHC-P

Recommended dilution

WB: 1:1000-1:5000

IHC-P: 1:50-1:100

Calculated MW

19 kDa

Observed MW

27 kDa

Host species

Rabbit


Clonality

Monoclonal


Clonality No.

DGR14242

Isotype	IgG
Purity	Affinity Purification
Conjugation	Un-conjugated
Storage Stability	Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.



Western blot analysis of extracts from K562, CHO-K1, HeLa cells using db11608 at 1:1000.



Immunohistochemical analysis of paraffin-embedded human breast cancer using db11608 antibody.