







DDIT3 (DGR14242) Rabbit mAb

db11608 Package : 10μL 20μL 50μL 100μL

Product Name: DDIT3 (DGR14242) Rabbit mAb

Cat.No.: db11608

Synonyms: CHOP; CEBPZ; CHOP10; CHOP-10; GADD153; C/EBPzeta

Application: WB, IHC-P

Reactivity: Human, Mouse, Rat

Host species: Rabbit

Background This gene encodes a member of the CCAAT/enhancer-binding protein (C/EBP) family of

transcription factors. The protein functions as a dominant-negative inhibitor by forming

heterodimers with other C/EBP members, such as C/EBP and LAP (liver activator protein), and preventing their DNA binding activity. The protein is implicated in adipogenesis and erythropoiesis,

is activated by endoplasmic reticulum stress, and promotes apoptosis. Fusion of this gene and

FUS on chromosome 16 or EWSR1 on chromosome 22 induced by translocation generates

chimeric proteins in myxoid liposarcomas or Ewing sarcoma. Multiple alternatively spliced

transcript variants encoding two isoforms with different length have been identified. [provided by

RefSeq, Aug 2010]

Immunogen A synthetic peptide of human DDIT3

Gene ID 1649

Swiss Prot P35638

Synonyms CHOP; CEBPZ; CHOP10; CHOP-10; GADD153; C/EBPzeta

Reactivity Human, Mouse, Rat

Application WB, IHC-P

Recommended dilution WB: 1:1000-1:5000

IHC-P: 1:50-1:100

Calculated MW 19 kDa

Observed MW 27 kDa

Host species Rabbit

Clonality Monoclonal

Clonality No. DGR14242






Isotype IgG

Purity Affinity Purification


Conjugation Un-conjugated

Storage Stability Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium


azide and 0.05% BSA. Stable for 12 months from date of receipt.



Western blot detection of DDIT3 in K562,CHO-K1,Hela cell lysates using DDIT3 antibody(1:1000 diluted).



Western blot analysis of extracts from HepG2, HCT116 cells using db11608 at 1:1000.



Immunohistochemical analysis of paraffin-embedded human breast cancer using db11608 antibody.