

DGRmAb® Recombinant

Securin (DGR16413) Rabbit mAb

db11881

Package : 10µL 20µL 50µL 100µL


Product Name : Securin (DGR16413) Rabbit mAb Cat.No.: db11881 Synonyms : EAP1; PTTG; HPTTG; TUTR1 Application : WB, IHC-P, ICC/IF, FC Reactivity : Human Host species : Rabbit


## Background

| Background           | The encoded protein is a homolog of yeast securin proteins, which prevent separins from                   |
|----------------------|---|
|                      | promoting sister chromatid separation. It is an anaphase-promoting complex (APC) substrate that           |
|                      | associates with a separin until activation of the APC. The gene product has transforming activity in      |
|                      | vitro and tumorigenic activity in vivo, and the gene is highly expressed in various tumors. The gene      |
|                      | product contains 2 PXXP motifs, which are required for its transforming and tumorigenic activities,       |
|                      | as well as for its stimulation of basic fibroblast growth factor expression. It also contains a           |
|                      | destruction box (D box) that is required for its degradation by the APC. The acidic C-terminal            |
|                      | region of the encoded protein can act as a transactivation domain. The gene product is mainly a           |
|                      | cytosolic protein, although it partially localizes in the nucleus. Three transcript variants encoding the |
|                      | same protein have been found for this gene. [provided by RefSeq, Sep 2013]                                |
| Immunogen            | A synthetic peptide of human Securin  |
| Gene ID              | 9232  |
| Swiss Prot           | O95997  |
| Synonyms             | EAP1; PTTG; HPTTG; TUTR1  |
| Reactivity           | Human   |
| Application          | WB, IHC-P, ICC/IF, FC   |
| Recommended dilution | WB: 1:2000-1:20000  |
|                      | IHC-P: 1:100-1:200  |
|                      | ICC/IF: 1:50  |
|                      | FC: 1:20-1:50   |
| Calculated MW        | 22 kDa  |
| Observed MW          | 25 kDa  |
| Host species         | Rabbit  |

## dvagbvo 戴格生物

| Clonality                | Monoclonal  |
|--------------------------|---|
| Clonality No.            | DGR16413  |
| lsotype                  | lgG   |
| Purity                   | Affinity Purification   |
| Conjugation              | Un-conjugated   |
| Storage Stability        | Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt. |
| 、<br>180-<br>130-<br>95- | Western blot detection of Securin in K562,3T3,Hela cell lysates using Securin antibody(1:1000 diluted).   |





Immunohistochemical analysis of paraffin-embedded human tonsil using db11881 antibody.