

Recombinant

DGRmAb®

**Histone H2A.X (DGR16840) Rabbit mAb**

db11980

Package : 10µL 20µL 50µL 100µL

**Product Name** : Histone H2A.X (DGR16840) Rabbit mAb**Cat.No.:** db11980**Synonyms** : H2AX; H2A.X; H2A/X**Application** : WB, IHC, ICC/IF, FC, IP**Reactivity** : Human, Mouse, Rat**Host species** : Rabbit**Background**

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene encodes a replication-independent histone that is a member of the histone H2A family, and generates two transcripts through the use of the conserved stem-loop termination motif, and the polyA addition motif. [provided by RefSeq, Oct 2015]

**Immunogen**

A synthetic peptide of human Histone H2A.X

**Gene ID**

3014

**Swiss Prot**

P16104

**Synonyms**

H2AX; H2A.X; H2A/X

**Reactivity**

Human, Mouse, Rat

**Application**

WB, IHC, ICC/IF, FC, IP

**Recommended dilution**

WB: 1:1000-1:5000

IHC: 1:200-1:1000

ICC/IF: 1:200-1:1000

FC: 1:20

IP: 1:20-1:100

**Calculated MW**

15 kDa


**Observed MW**

15 kDa


**Host species**

Rabbit


<b>Clonality</b>	Monoclonal
<b>Clonality No.</b>	DGR16840
<b>Isotype</b>	IgG
<b>Purity</b>	Affinity Purification
<b>Conjugation</b>	Un-conjugated
<b>Storage Stability</b>	Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.




Western blot analysis of extracts from 3T3, HeLa cells using db11980 at 1:1000.



Immunohistochemical analysis of paraffin-embedded human colon cancer using db11980 antibody.



Immunofluorescent analysis of HL-60 cells using db11980 antibody (green), and DAPI (blue).



Western blot analysis of extracts from PC-12 cells using db11980 at 1:1000.