Recombinant DGRmAb®


CDK6 (DGR11918) Rabbit mAb

db12115 **Package : 10μL 20μL 50μL 100μL**


Product Name : CDK6 (DGR11918) Rabbit mAb
Cat.No.: db12115
Synonyms : MCPH12; PLSTIRE
Application : WB, IHC-P, ICC/IF, FC
Reactivity : Human
Host species : Rabbit

Background	The protein encoded by this gene is a member of the CMGC family of serine/threonine protein kinases. This kinase is a catalytic subunit of the protein kinase complex that is important for cell cycle G1 phase progression and G1/S transition. The activity of this kinase first appears in mid-G1 phase, which is controlled by the regulatory subunits including D-type cyclins and members of INK4 family of CDK inhibitors. This kinase, as well as CDK4, has been shown to phosphorylate, and thus regulate the activity of, tumor suppressor protein Rb. Altered expression of this gene has been observed in multiple human cancers. A mutation in this gene resulting in reduced cell proliferation, and impaired cell motility and polarity, and has been identified in patients with primary microcephaly. [provided by RefSeq, Aug 2017]
Immunogen	A synthetic peptide of human Cdk6
Gene ID	1021
Swiss Prot	Q00534
Synonyms	MCPH12; PLSTIRE
Reactivity	Human
Application	WB, IHC-P, ICC/IF, FC
Recommended dilution	WB: 1:5000-1:50000 IHC-P: 1:100-1:200 ICC/IF: 1:200-1:500 FC: 1:100-1:1000
Calculated MW	37 kDa
Observed MW	37 kDa
Host species	Rabbit

Clonality	Monoclonal
Clonality No.	DGR11918
Isotype	IgG
Purity	Affinity Purification
Conjugation	Un-conjugated
Storage Stability	Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.



Western blot analysis of extracts from K562, HeLa cells using db12115 at 1:1000.



Immunofluorescence analysis of HeLa cells labelling CDK6 with db12115.

The cells were fixed with 4% PFA (10min, RT) followed by treatment with 0.1% Triton X-100 (10min, RT), and blocked in 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween 20 for 1h. The cells were then incubate with db12115 (1:200) at room temprature for 1h, followed by a further incubation at room temperature for 45min with Goat Anti Rabbit IgG (H+L)-AF488 (db10005, shown in green). Nuclear DNA was labeled in blue with DAPI.

Control: Secondary antibody only.