

CCL23 Rabbit pAb

db20454

Package : 20µL 50µL 100µL

Product Name : CCL23 Rabbit pAb**Cat.No.:** db20454**Synonyms** : CKb8; MIP3; Ckb-8; MIP-3; MPIF-1; SCYA23; Ckb-8-1; hmrp-2a; CK-BETA-8**Application** : WB, IHC**Reactivity** : Human**Host species** : Rabbit**Background**

This gene is one of several chemokine genes clustered on the q-arm of chromosome 17. Chemokines form a superfamily of secreted proteins involved in immunoregulatory and inflammatory processes. The superfamily is divided into four subfamilies based on the arrangement of the N-terminal cysteine residues of the mature peptide. This chemokine, a member of the CC subfamily, displays chemotactic activity on resting T lymphocytes and monocytes, lower activity on neutrophils and no activity on activated T lymphocytes. The protein is also a strong suppressor of colony formation by a multipotential hematopoietic progenitor cell line. In addition, the product of this gene is a potent agonist of the chemokine (C-C motif) receptor 1. Alternative splicing results in multiple transcript variants that encode different isoforms. [provided by RefSeq, Jul 2013]

Immunogen

A synthetic peptide of human CCL23

Gene ID

6368

Swiss Prot

P55773

Synonyms

CKb8; MIP3; Ckb-8; MIP-3; MPIF-1; SCYA23; Ckb-8-1; hmrp-2a; CK-BETA-8

Reactivity

Human

Application

WB, IHC

Recommended dilution

WB: 1:1000

IHC: 1:20

Calculated MW

13 kDa

Observed MW

13 kDa

Host species

Rabbit

Clonality

Polyclonal

Isotype

IgG

Purity	Affinity Purification
Conjugation	Un-conjugated
Storage Stability	Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.