

## CD43 Rabbit pAb

db4890

Package : 20µL 50µL 100µL

**Product Name** : CD43 Rabbit pAb**Cat.No.:** db4890**Synonyms** : LSN; CD43; GALGP; GPL115; LEU-22**Application** : WB, IHC, ICC/IF, FC**Reactivity** : Human**Host species** : Rabbit**Background**

This gene encodes a highly sialylated glycoprotein that functions in antigen-specific activation of T cells, and is found on the surface of thymocytes, T lymphocytes, monocytes, granulocytes, and some B lymphocytes. It contains a mucin-like extracellular domain, a transmembrane region and a carboxy-terminal intracellular region. The extracellular domain has a high proportion of serine and threonine residues, allowing extensive O-glycosylation, and has one potential N-glycosylation site, while the carboxy-terminal region has potential phosphorylation sites that may mediate transduction of activation signals. Different glycoforms of this protein have been described. In stimulated immune cells, proteolytic cleavage of the extracellular domain occurs in some cell types, releasing a soluble extracellular fragment. Defects in expression of this gene are associated with Wiskott-Aldrich syndrome. [provided by RefSeq, Sep 2017]

**Immunogen**

Recombinant protein of human CD43

**Gene ID**

6693

**Swiss Prot**

P16150

**Synonyms**

LSN; CD43; GALGP; GPL115; LEU-22

**Reactivity**

Human

**Application**

WB, IHC, ICC/IF, FC

**Recommended dilution**

WB: 1:1000

IHC: 1:500

ICC/IF: 1:20

FC: 1:20

**Calculated MW**

40 kDa

**Observed MW**

115 kDa

**Host species**

Rabbit

<b>Clonality</b>	Polyclonal
<b>Isotype</b>	IgG
<b>Purity</b>	Affinity Purification
<b>Conjugation</b>	Un-conjugated
<b>Storage Stability</b>	Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.