

TGF beta 1/3 Rabbit pAb

db527

Package : 20μL 50μL 100μL

Product Name : TGF beta 1/3 Rabbit pAb**Cat.No.:** db527**Synonyms** : CED; LAP; DPD1; TGFB; TGFbeta**Application** : WB**Reactivity** : Human, Mouse, Rat**Host species** : Rabbit**Background**

This gene encodes a secreted ligand of the TGF-beta (transforming growth factor-beta) superfamily of proteins. Ligands of this family bind various TGF-beta receptors leading to recruitment and activation of SMAD family transcription factors that regulate gene expression. The encoded preproprotein is proteolytically processed to generate a latency-associated peptide (LAP) and a mature peptide, and is found in either a latent form composed of a mature peptide homodimer, a LAP homodimer, and a latent TGF-beta binding protein, or in an active form consisting solely of the mature peptide homodimer. The mature peptide may also form heterodimers with other TGFB family members. This encoded protein regulates cell proliferation, differentiation and growth, and can modulate expression and activation of other growth factors including interferon gamma and tumor necrosis factor alpha. This gene is frequently upregulated in tumor cells, and mutations in this gene result in Camurati-Engelmann disease. [provided by RefSeq, Aug 2016]

Immunogen

Recombinant protein of human TGF beta 1

Gene ID

7040, 7043

Swiss Prot

P01137, P10600

Synonyms

CED; LAP; DPD1; TGFB; TGFbeta

Reactivity

Human, Mouse, Rat

Application

WB

Recommended dilution

WB: 1:1000

Calculated MW

44,47 kDa

Observed MW

44,12 kDa

Host species

Rabbit

Clonality

Polyclonal

Isotype	IgG
Purity	Affinity Purification
Conjugation	Un-conjugated
Storage Stability	Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.