

Recombinant

DGRmAb®

NFkB p105/p50 (DGR13934) Rabbit mAb

db12511

Package : 10µL 20µL 50µL 100µL

Product Name : NFkB p105/p50 (DGR13934) Rabbit mAb**Cat.No.:** db12511**Synonyms** : p50; KBF1; p105; EBP-1; CVID12; NF-kB1; NFkB-p50; NFkappaB; NF-kappaB; NFkB-p105; NF-kappa-B**Application** : WB, IHC-P**Reactivity** : Human,Mouse,Rat**Host species** : Rabbit**Background**

This gene encodes a 105 kD protein which can undergo cotranslational processing by the 26S proteasome to produce a 50 kD protein. The 105 kD protein is a Rel protein-specific transcription inhibitor and the 50 kD protein is a DNA binding subunit of the NF-kappa-B (NFkB) protein complex. NFkB is a transcription regulator that is activated by various intra- and extra-cellular stimuli such as cytokines, oxidant-free radicals, ultraviolet irradiation, and bacterial or viral products. Activated NFkB translocates into the nucleus and stimulates the expression of genes involved in a wide variety of biological functions. Inappropriate activation of NFkB has been associated with a number of inflammatory diseases while persistent inhibition of NFkB leads to inappropriate immune cell development or delayed cell growth. Alternative splicing results in multiple transcript variants encoding different isoforms, at least one of which is proteolytically processed. [provided by RefSeq, Feb 2016]

Immunogen

A synthetic peptide of human NFkB p105/p50

Gene ID

4790, 18033, 81736

Swiss Prot

P19838, P25799, Q63369

Synonyms

p50; KBF1; p105; EBP-1; CVID12; NF-kB1; NFkB-p50; NFkappaB; NF-kappaB; NFkB-p105; NF-kappa-B

Reactivity

Human,Mouse,Rat

Application

WB, IHC-P

Recommended dilution

WB: 1:1000-1:5000

IHC-P: 1:200-1:500


Calculated MW

105 kDa

Observed MW

105,50 kDa

Host species	Rabbit
Clonality	Monoclonal
Clonality No.	DGR13934
Isotype	IgG
Purity	Affinity Purification
Conjugation	Un-conjugated
Storage Stability	Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.



Western blot analysis of extracts from PC-12 cells using db12511 at 1:1000.