Recombinant DGRmAb®


NFkB p105/p50 (DGR13934) Rabbit mAb

db12511 **Package : 10μL 20μL 50μL 100μL**

Product Name : NFkB p105/p50 (DGR13934) Rabbit mAb
Cat.No.: db12511
Synonyms : p50; KBF1; p105; EBP-1; CVID12; NF-kB1; NFKB-p50; NFkappaB; NF-kappaB; NFKB-p105; NF-kappa-B
Application : WB, IHC-P
Reactivity : Human,Mouse,Rat
Host species : Rabbit

Background	This gene encodes a 105 kD protein which can undergo cotranslational processing by the 26S proteasome to produce a 50 kD protein. The 105 kD protein is a Rel protein-specific transcription inhibitor and the 50 kD protein is a DNA binding subunit of the NF-kappa-B (NFKB) protein complex. NFKB is a transcription regulator that is activated by various intra- and extra-cellular stimuli such as cytokines, oxidant-free radicals, ultraviolet irradiation, and bacterial or viral products. Activated NFKB translocates into the nucleus and stimulates the expression of genes involved in a wide variety of biological functions. Inappropriate activation of NFKB has been associated with a number of inflammatory diseases while persistent inhibition of NFKB leads to inappropriate immune cell development or delayed cell growth. Alternative splicing results in multiple transcript variants encoding different isoforms, at least one of which is proteolytically processed. [provided by RefSeq, Feb 2016]
Immunogen	A synthetic peptide of human NFkB p105/p50
Gene ID	4790, 18033, 81736
Swiss Prot	P19838, P25799, Q63369
Synonyms	p50; KBF1; p105; EBP-1; CVID12; NF-kB1; NFKB-p50; NFkappaB; NF-kappaB; NFKB-p105; NF-kappa-B
Reactivity	Human,Mouse,Rat
Application	WB, IHC-P
Recommended dilution	WB: 1:1000-1:5000 IHC-P: 1:200-1:500
Calculated MW	105 kDa
Observed MW	105,50 kDa

Host species	Rabbit
Clonality	Monoclonal
Clonality No.	DGR13934
Isotype	IgG
Purity	Affinity Purification
Conjugation	Un-conjugated
Storage Stability	Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.



Western blot analysis of extracts from PC-12 cells using db12511 at 1:1000.